

A Contribution to the Phenomenological Theory of Kinetic Processes in Ferromagnetic Dielectrics. I. The Relaxation in the Gas of the Spin Waves SOV/56-34-6-30/51

Akhiyezer, L. D. Landau and I. M. Lifshits for useful advice and also V. G. Bar'yakhtar and S. V. Peletminskiy for profitable discussions. There are 9 references, 6 of which are Soviet.

ASSOCIATION: Fiziko-tekhnicheskiy institut Akademii nauk Ukrainskoy SSR (Physics and Technical Institute, AS UkrSSR)

SUBMITTED: March 22, 1958

Card 3/3

NOV 56-35-2-23/60

24161

AUTHORS:

Vaganov, M. I., Tsukernik, V. M.

TITLE:

The Influence of Thermoelectric Forces on the Skin Effect in Metal (Vliyaniye termoelektricheskikh sil na skan-effekt v metalle)

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1958, Vol 35, Nr 2, pp 474-478 (USSR)

ABSTRACT:

Formulae are deduced for the surface resistance in consideration of thermoelectric forces. The following system of equations serves as a basis:

$$\text{curl } \vec{H} = \frac{4}{c} \vec{j}; \text{curl } \vec{E} = -\frac{1}{c} \frac{d\vec{H}}{dt}; c \frac{\partial \theta}{\partial t} + \text{div } \vec{q} = 0;$$

$$\vec{E}_i = \rho_{ik} j_k + \alpha_{ik} \partial \theta / \partial x_k; q_i = T \alpha_k j_k - \chi_{ik} \partial \theta / \partial x_k$$

where θ is the high-frequency addition to the average temperature T of the sample, c - the specific heat of the metal, \vec{q} the heat current, χ_{ik} - the resistance tensor, ρ_{ik} - the thermal conductivity tensor, and α_{ik} - the tensor of the thermoelectrical coefficients. ($\alpha_{ik} = \alpha_{ki}$).

Card 1/2

SCV/86-34-2-23/60

The Influence of Thermoelectric Forces on the Skin Effect in Metal

For the surface impedance it holds that $\xi_x = E_x(0)/H_y(0)$ and $\xi_y = -E_y(0)/H_x(0)$; $\xi_x = \xi_{gx} = \sqrt{\omega \sigma_{xx}/4\pi i} = \sqrt{\omega \sigma_1/4\pi i}$; ($\sigma_{11} = \sigma_{22} = \sigma_{33}$; $\sigma_{11} = \sigma_{11}$) and ξ_y is decomposed into ξ_y^{ad} and ξ_y^{isoth} , for which equations are deduced as well. In the case of isotropy $\xi = 1/\sqrt{\epsilon - \sin^2 \phi}$ and $\xi^{isoth} = \sqrt{\epsilon - \sin^2 \phi}/\epsilon$ is obtained, and for ξ^{ad} a considerably more complicated expression is found, the first term of which is equal to the expression for ξ^{isoth} . In conclusion, the authors express their gratitude to L. D. Landau for his valuable discussions. There are 2 references, 2 of which are Soviet.

ASSOCIATION: Fiziko-tekhnicheskiy institut Akademii nauk ^USSR (Physico-Technical Institute, ~~AS~~ USSR)

SUBMITTED: March 26, 1958
Card 2/2

24(3)
AUTHORS:

Kaganov, M. I., Tsukernik, V. M.

SOV/56-36-1-30/62

TITLE:

On the Phenomenological Theory of the Kinetic Processes
in Ferromagnetic Dielectrics (K fenomenologicheskoy teorii
kineticheskikh protsessov v ferromagnitnykh dielektrikakh)
II. The Interaction of Spin Waves With Phonons
(II. Vzaimodeystviye spinovykh voln s fononami)

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1959,
Vol 36, Nr 1, pp 224-232 (USSR)

ABSTRACT:

The spin waves have to be described by the vector of the
density of the magnetic moment $\vec{M} = \vec{M}(\vec{r}, t)$ by the deformation
tensor $u_{ik} = (1/2) (\partial u_i / \partial x_k + \partial u_k / \partial x_i)$ where $u = u(\vec{r}, t)$
denotes the vector of displacement. In this case, the interaction
Hamiltonian will be the sum of all components in the expansion
of the energy of the ferromagnetic with respect to the powers
of M_i , $\partial M_i / \partial x_k$, and u_{ik} : $\gamma_{iklm} \int M_i M_k u_{lm} dv$,

$\lambda_{iklmrs} = \int \frac{\partial M_i}{\partial x_k} \frac{\partial M_l}{\partial x_m} u_{rs} dv$. The first of these terms corresponds

Card 1/3

to the energy of magnetostriction. The interaction Hamiltonian

On the Phenomenological Theory of the Kinetic Processes in Ferromagnetic Dielectrics. II. The Interaction of Spin Waves With Phonons

SOV/56-56-1-30/52

is written down as $H_{\text{interaction}} = H_{\text{magnetostriction}} + H_{\text{exchange}}$,
where $H_{\text{interaction}} = \gamma \int M_i M_k u_{ik} dv$,

$$H_{\text{exchange}} = \lambda_1 \int \frac{\partial M_1}{\partial x_i} \frac{\partial M_1}{\partial x_k} u_{ik} dv + \lambda_2 \int \frac{\partial M_1}{\partial x_i} \frac{\partial M_1}{\partial x_i} u_{kk} dv.$$

In the second part of the paper the production and the absorption of phonons by spin waves are calculated. The authors also investigate the energy transfer from spin waves to phonons. The spin waves and the phonons are described by equilibrium distribution functions of the Bose (Boze) type for different temperatures. The coefficient of heat transfer for the activation process of the emission and absorption of phonons decreases exponentially with temperature. The third part of the present paper deals with the annihilation of 2 spin waves with production of a phonon and with the decay of a phonon into 2 spin waves. These processes are due to the magnetostriction part of the Hamiltonian. Practically at any temperature above μMo , production and absorption of phonons

Card 2/3

On the Phenomenological Theory of the Kinetic
Processes in Ferromagnetic Dielectrics. II. The Interaction
of Spin Waves With Phonons

SOV/56-36-1-30/62

play the most important part among the one-phonon processes. The two-photon processes are described by the square terms of the expansion of the energy of the ferromagnetic with respect to the powers of the deformation tensor. There are 2 types of such processes: scattering of a phonon on a spin wave (or vice versa) and emission (or absorption) of 2 phonons by a spin wave. Finally, the time necessary for the establishment of equilibrium (relaxation time) between spin waves and phonons is calculated. There are 5 Soviet references.

ASSOCIATION: Fiziko-tekhnicheskiy institut Akademii nauk Ukrainskoy SSR
(Physico-Technical Institute of the Academy of Sciences,
Ukrainskaya SSR)

SUBMITTED: July 8, 1958

Card 3/3

24 (3)
AUTHORS:

Kaganov, M. I., Tsukernik, V. M.

SOV/56-37-3-35/62

TITLE:

Nonresonance Absorption of the Energy of a Variable Magnetic Field by a Ferromagnetic Dielectrics

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1959,
Vol 37, Nr 3 (9), pp 823-832 (USSR)

ABSTRACT:

A ferromagnetic substance may absorb the energy of a variable magnetic field in various ways: either the magnetic field (the amplitude of which is to be small, $H' \sim e^{-i\omega t}$) is polarized perpendicular to the equilibrium magnetic moment, in which case the magnetic field turns the moment without changing its value, or the field is polarized in the direction of the moment, in which case a superposed magnetization occurs, i.e. the absolute value of the magnetic moment is varied. In both cases dissipative processes are caused, which are interrelated with the interaction of spin waves, both among themselves, and also with phonons (it is assumed that the dielectric substance is magnetized up to saturation). In the present paper the authors confine their attention to those in which the variable magnetic field (frequency ω) is polarized in the direction of the

Card 1/4

Nonresonance Absorption of the Energy of a Variable
Magnetic Field by a Ferromagnetic Dielectrics

SOV/56-37-3-35/62

equilibrium moment. Calculation of absorption is then reduced to calculation of the imaginary part of longitudinal magnetic susceptibility $\mu_{||} = \mu' + i\mu''$. The complex character of the relaxation processes in ferromagnetic dielectrics leads to a complex frequency dependence $\mu_{||}(\omega)$. The authors proceed from the simple model, in which the ferroelectric body is considered to have a magnetic moment, the vibrations of which propagate in form of spin waves. This is justified at sufficiently low temperatures, when the vibrations of one of the magnetic sublattices are not excited by the others. The theory of spin waves may be applied to real bodies, e.g. to ferrites the semiconductor properties of which at low temperatures are insignificant (the number of free electrons with decreasing temperature tends exponentially towards zero). The first part of this paper deals with the low frequency range, i.e. ω is much smaller than the reciprocal spin-spin relaxation time τ_{ss} : $\omega\tau_{ss} \ll 1$. The following is obtained for the imaginary part of the longitudinal magnetic susceptibility after all frequency- and temperature ranges have been dealt with in detail:

Card 2/4

Nonresonance Absorption of the Energy of a Variable
Magnetic Field by a Ferromagnetic Dielectrics

SOV/56-37-3-35/62

$$\mu'' \sim \begin{cases} \frac{M_0 \mu_c}{\theta_c} \left(\frac{T}{\theta_c} \right)^{1/2} \frac{\omega \tau_{sl}}{1 + \omega^2 \tau_{sl}^2}, & \omega \sqrt{\tau_{sl} \tau_{ss}} \ll 1, \quad T \ll T_0 \\ \frac{\mu M_0}{\theta_c} \left(\frac{T}{\theta_c} \right)^{1/2} \frac{\omega \tau_{ss}}{1 + \omega^2 \tau_{ss}^2}, & \omega \sqrt{\tau_{sl} \tau_{ss}} \gg 1, \quad T \ll T_0 \\ \frac{\hbar \omega}{\mu M_0} \frac{1 + \omega^2 \tau_2^2}{(1 + \omega^2 \tau_1^2)^2 + \omega^2 \tau_3^2}, & \omega \tau_{ss} \ll 1, \quad T \gg T_0 \end{cases}$$

Here $T_0 \sim \mu M_0 (\theta_c / \mu M_0)^{3/7}$, τ_{sl} denotes the spin-lattice relaxation time. The second part deals with high frequencies ($\omega \gg 1/\tau_{ss}$), the Hamiltonian of the interaction between field and spin being set up according to Kaganov (Ref 2):

$\mathcal{H}_{int} = \mu \int \hat{H}_z \hat{a}^*(\vec{r}) \hat{a}(\vec{r}) d\vec{r}$. If ω tends towards ∞ , then magnetic susceptibility tends towards unity, and its imaginary part towards zero. For the frequency dependence of μ'' at very high

Card 3/4

Nonresonance Absorption of the Energy of a Variable
Magnetic Field by a Ferromagnetic Dielectrics

SOV/56-37-3-35/62

frequencies it holds that $\mu'' \approx \frac{8\sqrt{2}}{15} \frac{u^2}{a^3} \frac{(2\pi\mu H_0)^2}{\theta_c^{3/2} (H\omega)^{3/2}}$,

$(H\omega \gg \frac{h}{\tau_{ss}}, T, \mu H_{eff})$. Thus, μ'' decreases inversely proportionally with $\omega^{3/2}$ and is independent of temperature. The authors thank V. G. Eas'yakhtar for valuable discussions. There are 10 Soviet references.

ASSOCIATION: Fiziko-tekhnicheskiy institut Akademii nauk Ukrainskoy SSR
(Physico-technical Institute of the Academy of Sciences,
Ukrainskaya SSR)

SUBMITTED: April 18, 1959

Card 4/4

21(0)
ATTN:

Chernikov, I.

308/55-67-4-77

8797148:

The Fifth All-Union Conference on the Physics of Low Temperatures (5-y^e Vsesoyuznoye soobshcheniye po fizike nizkikh temperatur)

PHOTOGRAPH

Voprosy fizicheskikh nauk, 1959, Vol 67, No 4, pp 743-750
(USSR)

Abstract

This conference took place from October 27 to November 1 at Tallin. It was organized by the Ukrainian higher-school scientific staff Andreiev, rank SSSt (Department of Physico-mathematical sciences of the Academy of Sciences, USSR), the Academician rank Grynkozyy SSt (Academy of Sciences, Ukrainian SSR), and the Collectively Governmentally authorized in Stalin (Collective State University named Stalin). The Conference was attended by about 500 specialists from Tallin, Moscow, Leningrad, Kiev, Lvov, Kyiv, Zhytomyr, and other cities as well as by a number of young Chinese scientists working in the USSR. About 600 scientific articles, of which were divided according to research fields.

III. Galygona, the "Linnéan"
and the Pöschonoxly (Kistner)

Cher'kov Physico-technical Institute, Cher'kov University

showed that the most important part in connection with the electrocatalytic properties of metals is played by the concentration of the metal ions in the solution.

form of the Fermi surface of conductivity electrons. It was found that the concentration of the impurities can be carried out

Alkeevskiy (197) spoke about the various contributions made by the USSR to the development of the world economy.

of the resistance in the fundamental magnetic field at heat treatment of Au-Cu-Fe-3Ni and (together with

I. I. Kostina) of St. Ye. S. Borovik and V. G. Volbitsky

(KARYI) investigated the calorimetric properties of the mixtures of chromolite and zirconium and found that the

Resistance of chromium grows with field strength although the rate of increase is not linear and is dependent on the field strength.

indicating a recombination value of 2.0×10^{-4} for the recombination minimum in gold at low temperatures.

tures and found that if the sample is heated, the minimum T_c of P-Cardukov (IPP) laid in this connection

in the course of the discussion that the minicore effect does

not occur in gold in the case of very pure samples. The occurrence of the minimum is explained by the plastic deformation of the material.

tion of the sample at helium temperatures. Mo. In. Assoc.

(KAPL) gave a report of the results of the high-frequency resistance of steel in aquatics; theory of the high-frequency resistance of steel in aquatics; and the high-frequency resistance of steel in aquatics.

at a constant magnetic field at low temperatures. V. A. Kabanov and V. M. Tsukernik (KhFTI) spoke about a theoretical investigation of the

Effect of the Influence exerted by thermoelectric force
in various conductors. B. T. Yek'in and

upon the skin effect at various frequencies. Dr. V. Alexandrov (KhTI) spoke about measurements of the

electric resistance of thin wires. Data from nitrogen-pure aluminum and cadmium, and computed the free length of

path at 4.2°K in these solids amounting to $1/3$ to $2/3$ of the total path.

X. H. Brandt (MCU) and R. A. Seikaly and
(FERT) investigated the influence exercised by the hydro-

static pressure (or 1000 atmospheres absolute pressure) up to 1000 atmospheres and investigated

the quantum oscillations of the magnetic susceptibility of

March at 1.6 - 4.2°F. O. Ye. "Liberman and A. J. Rose
1967) gave a theoretical explanation of the facts in

already relatively small deformations exercise considerably

influence upon oscillation effects in nitrate. At the same time, A. I. Borovikh-Romanov (LPP) delivered a report on investigations

stions he carried out of the anisotropy of the weak ferro-

magnetism in monocrystals, compared to the effect of anisotropy was predicted by the theory of the effect of anisotropy on the conductivity of the crystal.

dynamical theory developed by Dykhovitskiy/. In the course of the discussion B. A. Alkhimov (YPP) spoke about neutro-

graphical investigations he carried out of the magnetic

structure of MnCO_3 and FeCO_3 as for MgCO_3 and CaCO_3 . The importance of the method based upon implicit stresses, the importance of the method based upon

Dyalozhinsky's theory. M. E. Feyzes (VIL'NIL), whose report was read by A. I. Borovik-Buzinov, reported on the

ments carried out by him (in the IPF) of the paginated and

trophy of the antiferromagnetic CuSO₄ and CoSO₄-monohydrate

Y. A. Yurov (171 A. 5334, SWERDLOV) above name

Card 7/11

Case# 6/11

Case# 5/11

V. M. TSOKERNIK

304/55-67-4-7/7

Chentsov, R.
The Fifth All-Union Conference on the Physics of Low
Temperatures (5-ye Vsesoyuznyye soveshchaniya po fizike nizkikh
temperatur)

24(0)
AUTHOR:
TITLE:

PERIODICAL:
(USSR)

ABSTRACT:
This Conference took place from October 27 to November 1 at
Tbilisi; it was organized by the Ondeleniye fiziko-matemati-
cheskikh nauk Akademii na SSSR (Department of Physico-
mathematical Sciences of the Academy of Sciences, USSR),
the Akademiyi nauk Gruzinskoy SSR (Academy of Sciences,
Gruzinskaya SSR), and the Tbilisskiy gosudarstvennyy uni-
versitet im. Sh. Rustaveli (Tbilisi State University imeni
Shalvaya Rustaveli). The Conference was attended by about 300 specialists from
the USSR, the GDR, Poland, Czechoslovakia, Rumania, and
Yugoslavia, as well as by a number of young scientists
studying in the USSR. About 50 lectures were deliv-
ered which were divided according to research fields.

IV. Magnetism.
A. S. Borovik-Romanov (IFF) delivered a report on investiga-
tions he carried out of the anisotropy of the weak ferro-
magnetism in monocystal samples of the antiferromagnetic
MnCO₃ (the effect of anisotropy was predicted by the thermo-
dynamical theory developed by A. I. Zhuravskiy). In the course
of the discussion A. I. Zhuravskiy (IFF) spoke about neutron-
graphical investigations he carried out of the magnetic
structure of MnCO₃ and FeCO₃ at low temperatures. P. L.
Kapitanov (IFF) stressed the importance of the method based upon
Kapitanyuk's theory. A. K. Kreyzas (VNIIFRIL), whose
report was read by A. S. Borovik-Romanov, reported on measur-
ements carried out by him (in the IFF) of the magnetic aniso-
tropy of the antiferromagnetic CuSO₄ and FeSO₄ monocystals.

Card 7/11

Ye. A. Furor (IFF AN SSSR, Sverdlovsk) spoke about his theo-
retical investigations of the magnetisability, the susceptibility,
the specific heat, and the resonance frequencies of anti-
ferromagnetics and ferrimagnetics. A. I. Sudovtsov and
Ye. A. Furor (IFF AN SSSR) spoke about measurements of the
electric resistance of iron in magnetic fields in a wide tem-
perature range with simultaneous plotting of the magnetization
curve. A. I. Zhuravskiy (IFF AN SSSR) spoke about measurements
of the Hall effect and the Hall effect of polycrystalline samples,
nickel and Fe-Ni at low temperatures. Ye. I. Kondratyuk

V. Roda, B. Gryn and Ch. Shchukun (IGU) gave a report
on susceptibility measurements on nickel and its alloys with
copper at low temperatures. A. I. Sandak (IGU) spoke about
measurements of the paramagnetic resonance of Fe³⁺ in terbium
nitrate at temperatures of liquid hydrogen. M. I. Kaganov
and V. M. Tsokernik (IFF) dealt with the kinetic phenomena
in ferromagnetics at low temperatures and with calculation
of relaxation times. A. I. Akhlyev, V. Sh. Zhuravskiy and
P. L. Kapitanov (IFF) carried out a theoretical investigation
of the relaxation of the magnetic moment in ferromagnetics.
V. M. Tsokernik (IFF AN SSSR) showed that a lightly polarized elastic
(ultrasonic) wave of a frequency of 10⁶ cycles when passing
through a ferromagnetic substance in the direction of the
magnetic field, is subjected to turn of the plane of polarization
of the order of 10⁻³ - 10⁻⁴ rad/cm oriented. M. I.
Kaganov pointed out that in this connection yet another
phenomenon may be observed, namely, the resonance absorption
of ultrasonics if the wavelength is equal to the radius of
the Larmor orbit of the electron. V. V. Varshniy (IFF AN SSSR)

Card 8/11

85974

S/126/60/010/005/029/030
E032/E414

24.2200 (1385,1144,1162)

AUTHORS: Kaganov, M.I., Tsukernik, V.M. and Chupis, I.Ye.
TITLE: Theory of Relaxation Processes in Antiferromagnetics ↑
PERIODICAL: Fizika metallov i metallovedeniye, 1960, Vol.10, No.5,
pp.797-798

TEXT: The method put forward by Akhiezer (Ref.1) and Kaganov and Tsukernik (Ref.2) is used to calculate the mean probabilities of processes associated with the interaction between spin waves in ferromagnetics. The theory holds in the temperature region

$$\omega_c \left(\frac{\mu_0 M_0}{\omega_c} \right)^{1/2} \ll T \ll \omega_c$$

in which the energy of the spin wave is given by the well-known expression

$$\epsilon_\lambda = \omega_c (ak_\lambda)$$

Card 1/3

85974

S/126/60/010/005/029/030
E032/E414

Theory of Relaxation Processes in Antiferromagnetics

The notation is defined in the previous paper by the present authors (Ref.3). The processes considered are: (a) combination of two spin waves into one, and (b) collision of two spin waves. The probability of other processes, for example combination of three spin waves into one, have zero probability since energy and momentum conservation laws cannot be satisfied at the same time. It is found that the mean probabilities for the above two processes are respectively given by

$$\bar{W}_3 \approx \frac{\omega_c}{h} \left(\frac{\mu_0 M_0}{\omega_c} \right)^{5/2} \frac{T}{\omega_c}^3 \quad (3)$$

$$\bar{W}_4 \approx \frac{\omega_c}{h} \left(\frac{T}{\omega_c} \right)^5 \quad (4)$$

Comparison of these two probabilities shows that in the above
Card 2/3

85974

S/126/60/010/005/029/030

E032/E414

Theory of Relaxation Processes in Antiferromagnetics

temperature region the non-homogeneous exchange interaction is responsible for setting up the thermodynamic equilibrium in a spin wave system. Acknowledgments are expressed to V.G.Bar'yakhtara for valuable discussions. There are 3 Soviet references.

ASSOCIATIONS: Fiziko-tehnicheskii institut AN USSR
(Physical-Technical Institute AS UkrSSR)
Khar'kovskiy gosudarstvennyy universitet
im. A.M.Gor'kogo (Khar'kov State University
im. A.M.Gor'kiy)

SUBMITTED: April 18, 1960

Card 3/3

83743

S/056/60/038/004/036/048
B006/B056

9.4300 (1035, 1138, 1143)

24.7900

AUTHORS:

Kaganov, M. I., Tsukernik, V. M.

TITLE:

Nonresonance Absorption of the Energy of an Alternating
Magnetic Field by a Ferromagnetic Dielectric. II

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1960,
Vol. 38, No. 4, pp. 1320 - 1325

TEXT: In part I of the present paper, the nonresonance absorption of the energy of an alternating magnetic field by a ferromagnetic dielectric has already been investigated on the assumption that this field is polarized in the direction of the axis of easiest magnetization. The case is now dealt with, in which the field is polarized perpendicular to this axis. These two cases are described as longitudinal and transverse fields. Contrary to what is the case in a longitudinal field, energy absorption may occur in a transverse field even if no dissipative processes occur, but when the field frequency agrees with the ferromagnetic resonance frequency. Neither resonance nor the effects connected with it are considered. The results obtained here relate to frequencies

Card 1/3

83743

Nonresonance Absorption of the Energy of an
Alternating Magnetic Field by a Ferro-
magnetic Dielectric. II

S/056/60/038/004/036/048
B006/B056

that are distant from resonance. The dissipative processes connected with the energy absorption are caused by the interaction between the magnetic field and the spin waves. The authors assume (like in Ref. 1) that the ferromagnetic dielectric is magnetized up to saturation at a given temperature, i.e., that it may be considered to be a single-domain sample. The sample is also considered to be sufficiently pure, so that impurity effects may be neglected. It is shown that the nonresonance absorption of magnetic field energy is connected with the forming of two spin waves as a result of a "collision" between a photon and a spin wave. Besides, an absorption of the photon in higher perturbation-theoretical approximations is possible due to processes in which a large number of spin waves participate. The most important process in this case is one of the fifth order, which occurs in second perturbation-theoretical approximation. The coefficient of the absorption of magnetic field energy $\Gamma = Q \left(\frac{h\omega}{8\pi} V \right)^{-1}$ is now investigated, where Q denotes the amount of energy absorbed per second within the entire sample volume. X

Card 2/3

83743

Nonresonance Absorption of the Energy of an Alternating Magnetic Field by a Ferromagnetic Dielectric. II

S/056/60/038/004/036/048
B006/B056

One obtains $\Gamma = \frac{1}{32\pi^2} \frac{\omega^2 T^2}{\mu M_0 \theta_c^3} \omega (1 - e^{-\hbar\omega/T}) I(\eta, \nu)$, where $\eta = \mu H_e/T$ and

$\nu = \hbar\omega/T$; the function $I(\eta, \nu)$ is represented as an integral equation by (13) and (13'). These equations are integrated for the special cases of low frequencies ($\nu \ll 1, \eta$), resonance frequency ($\nu = \eta$), and high frequencies ($1 \gg \nu \gg \eta$; $\nu \gg 1 \gg \eta$; $\eta \gg \nu \gg 1$; $\nu \gg \eta \gg 1$). The resulting asymptotic values of the integrals are used to determine approximate formulas for Γ in the various special cases. The authors thank A. I. Akhiezer and V. G. Bar'yakhtar for discussions. There are 3 references: 2 Soviet and 1 US.

ASSOCIATION: Fiziko-tekhnicheskiy institut Akademii nauk Ukrainskoy SSR
(Institute of Physics and Technology of the Academy of Sciences Ukrainskaya SSR)

SUBMITTED: November 23, 1959

Card 3/3

S/056/60/039/002/044/044
B006/B070

AUTHORS: Kaganov, M. I., Tsukernik, V. M.

TITLE: Theory of the Non-resonant Absorption of a Variable
Magnetic Field by a Ferromagnetic Dielectric ¹

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1960,
Vol. 39, No. 2(8), p. 518

TEXT: In an earlier paper, the authors calculated the absorption coefficient of a variable magnetic field polarized perpendicular to the axis of easiest magnetization. The Hamiltonian used in that case took only the exchange interaction into account. Later, Tsukernik showed that in this case only resonance absorption of a homogeneous magnetic alternating field may take place, since the total magnetic moment commutes with the Hamiltonian of the system; the results of the earlier paper (Ref. 1) are therefore wrong. The error is related to the fact that in the calculation of the matrix elements of the transition, the authors restricted themselves to the first approximation of the perturbation theory. A consideration of the second approximation shows that the matrix

Card 1/2

Theory of the Non-resonant Absorption of a
Variable Magnetic Field by a Ferromagnetic
Dielectric

S/056/60/039/002/044/044
B006/B070

element vanishes to the corresponding accuracy. The non-resonant absorption of the homogeneous field is related to the relativistic interactions in the system (dipole-dipole interaction, anisotropy energy, etc.). These problems will be considered in a later paper. Here, the authors give only a formula for the coefficient of absorption of a transverse magnetic field whose frequency is considerably smaller than that of the spin wave in the case of a zero quasi-momentum. The absorption described by this formula is due to a decay of a photon into two spin waves with oppositely directed quasi-momenta. There is 1 Soviet reference. ✓

ASSOCIATION: Fiziko-tekhnicheskiy institut Akademii nauk Ukrainskoy SSR
(Institute of Physics and Technology of the Academy of
Sciences Ukrainskaya SSR)

SUBMITTED: July 2, 1960

Card 2/2

ACCESSION NR: APL019844

S/0181/64/006/003/0818/0826

AUTHORS: Genkin, G. M.; Golubeva, N. G.; Tsukornik, V. M.

TITLE: The spin phonon width of lines of antiferromagnetic resonance

SOURCE: Fizika tverdogo tela, v. 6, no. 3, 1964, 818-826

TOPIC TAGS: spin phonon interaction, magnetic resonance, magnetization precession, exchange interaction, antiferromagnetism

ABSTRACT: The authors have examined the relaxation of the uniform precession of magnetization in a uniaxial antiferromagnetic as a consequence of interaction between spin waves and phonons (both acoustical and optical). They show that the relaxation processes of creating (or absorbing) an acoustical phonon by a spin wave, by virtue of the law of conservation of energy and momentum, may take place only at frequencies of

$$\omega_0 > \omega_0^{\pm} = \frac{\sqrt{3}h_A v}{a}$$

where ω_0 is the frequency of antiferromagnetic resonance, h_A is the field of

Card 1/2

ACCESSION NR: AP4019844

anisotropy, v the velocity of sound, and a the lattice constant. The interaction is an exchange process. The width of the line has been computed for any temperature. At rather high frequencies of antiferromagnetic resonance, $\omega_0 > \omega_0^{gr}$, where normally $\omega_0^{gr} \sim 2 \cdot 10^{12}$ sec⁻¹deg, the spin-phonon line width at low temperatures proves to be much greater than spin-spin line width. At frequencies of $\omega_0 < \omega_0^{gr}$, the spin-phonon width is small compared to spin-spin width, except for regions of very low temperatures ($T < 10^{-3}$ K). Orig. art. has: 39 formulas.

ASSOCIATION: Radiofizicheskiy institut, Gorkiy (Radiophysical Institute)

SUBMITTED: 21Sep63

DATE ACQ: 31Mar64

ENCL: 00

SUB CODE: SS, EM

NO REF SOV: 003

OTHER: 008

Card 2/2

24.7900 1147 1158 1163, 1144

30083
S/048/61/025/C11/007/031
B108/B138

AUTHORS: Kaganov, M. I., and Tsukernik, V. M.

TITLE: Off-resonance absorption of a variable magnetic field by ferromagnetic dielectrics at low temperatures

PERIODICAL: Akademiya nauk SSSR. Izvestiya. Seriya fizicheskaya, v. 25, no. 11, 1961, 1346-1351

TEXT: The authors studied the absorption of an h.f. magnetic field whose plane of circular polarization is perpendicular to the equilibrium magnetic moment of the ferromagnetic. They consider absorption without resonance, which can only occur by the interaction of spin waves with one another and with phonons. The calculations are made for temperatures considerably below the Curie point θ_c . The absorption coefficient Γ of the magnetic field is given as $\Gamma = \frac{8\pi k_B}{h^2 V} \sum \{w_{if} - w_{fi}\}$, where w_{if} is the probability of direct, and w_{fi} of inverse, transition. The absorption

Card 1/4

30063 S/048/61/025/011/007/031
B108/B138

Off-resonance absorption of a...

coefficient due to spin-spin interaction is evaluated for two limiting cases: (1) High temperatures $T \gg \varepsilon_0$ (but still $T \ll \theta_c$):

$$\Gamma_{ss} \approx \begin{cases} A_1 \omega \left(\frac{w}{\theta_c} \right)^2 \frac{\mu M_1}{\theta_c} \cdot \frac{\hbar \omega T^2}{\varepsilon_0^3}, & \hbar \omega \ll \varepsilon_0, \\ A_2 g M_0 \left(\frac{w}{\theta_c} \right)^2 \frac{T^2}{\hbar \omega \theta_c}, & \varepsilon_0 \ll \hbar \omega \ll T, \\ \frac{4}{15} \sqrt{\frac{\pi}{2}} \zeta(3/2) g M_0 \left(\frac{w}{\theta_c} \right)^2 \frac{T^{3/2}}{(\hbar \omega)^{1/2} \theta_c}, & \hbar \omega \gg T, \end{cases} \quad (17)$$

where $w = \frac{\mu^2}{R^3}$ is the dipole-dipole interaction energy; $A_1 \sim A_2 \sim 10^3$.

(2) Low temperatures ($T \ll \varepsilon_0$):

Card 2/4

Off-resonance absorption of ...

30953, 048/61/025/011/007/031
B100/B136

$$\Gamma_{ss} \approx \begin{cases} B_1 \omega \left(\frac{\omega}{\theta_c} \right)^2 \frac{\mu M_0}{\theta_c} \frac{\hbar \omega T^{1/2}}{\varepsilon_0^{1/2}} e^{-\frac{\pi \varepsilon_0}{T}}, & \hbar \omega \leq \varepsilon_0, \\ \frac{4}{15} \sqrt{\frac{\pi}{2}} \omega \left(\frac{\omega}{\theta_c} \right)^2 \frac{\mu M_0}{\theta_c} \left(\frac{T}{\varepsilon_0} \right)^{1/2} e^{-\frac{\pi \varepsilon_0}{T}}, & \hbar \omega \geq \varepsilon_0, \end{cases} \quad (18)$$

where

$$B_1 = \frac{15\pi^{1/2}}{8\sqrt{2}} \left\{ 121 + \frac{1}{15} \left[\left(\frac{2\beta}{\pi} - \frac{2}{3} \right)^2 + \frac{121}{45} \right] \right\} + \frac{4}{15} \sqrt{\frac{\pi}{2}} \approx 10^4$$

$\varepsilon_0 = \mu H_0 + \beta \mu M_0$ where H_0 denotes the permanent magnetic field along the axis of easiest magnetization, M_0 the equilibrium magnetic moment, β the anisotropy constant, a the lattice constant, $\mu = g\hbar$, g the gyromagnetic ratio. The results for spin-lattice interaction show that these contributions to the overall absorption coefficient are considerably smaller than the contribution due to spin-spin interaction. The considerations have been made for perfect ferromagnetics without

Card 3/4

Off-resonance absorption of a...

3/048/61/025/011/007/031
B108/B138

impurities. I. M. Lifshits is thanked for discussions. This paper was read at the Conference on ferromagnetism and antiferromagnetism in Leningrad, May 5-11, 1961. There are 1 figure and 4 Soviet references.

ASSOCIATION: Fiziko-tekhnicheskiy institut Akademii nauk USSR
(Physicotechnical Institute of the Academy of Sciences
UkrSSR)

Card 4/4

24.7900 (1055, 1144, 1163)
24.2200 1164 1068 1121

300614
S/048/61/025/011/008/031
B108/B138

AUTHORS: Kaganov, M. I., and Tsukernik, V. M.

TITLE: An absorption mechanism for a longitudinal magnetic field by a ferromagnetic dielectric

PERIODICAL: Akademiya nauk SSSR. Izvestiya. Seriya fizicheskaya, v. 25, no. 11, 1961, 1352-1353

TEXT: The ferro-dielectric absorption of the energy of a variable magnetic field polarized along the axis of easiest magnetization was studied in detail in a previous paper (Ref. 1: Zh. eksperim. i teor. fiz., 37, 823 (1959)). The absorption of the field at $\omega\tau_{ss} \ll 1$, where τ_{ss} is relaxation time, was calculated with the aid of the kinetic equations for the distribution function of the spin waves. The calculations were made for temperatures T considerably below Curie point θ_c . At high frequencies, absorption is chiefly determined by the disintegration of a photon into two spin waves with opposite momenta. This process was also considered in the above paper. It was pointed out, however, that a photon will only

Card 1/4

30061
S/048/61/025/011/008/031
B108/B138

An absorption mechanism for a...

disintegrate into two spin waves at frequencies ω greater than $\frac{2\varepsilon_0}{\hbar}$, where ε_0 is the energy of a zero-momentum spin wave. The wide range $\frac{1}{\tau} \ll \omega < \frac{2\varepsilon_0}{\hbar}$ has not been considered as yet. Absorption in this range is the subject of the present paper. In the lowest spin-wave approximation, the Hamiltonian for spin wave interaction with one another and with a longitudinal magnetic field is $H' = H_{\text{int}} + H_h$ (1) where $H_{\text{int}} = \sum_{123} \Phi_{123} a_1^+ a_2^+ a_3 + \text{complex conjugate}$ (2) and $H_h = \frac{1}{2} \mu h_0 (e^{i\omega t} + e^{-i\omega t}) \sum_k a_k^+ a_k$ (2'). h_0 denotes the amplitude of

the variable field. The other notations may be found in Ref. 2 (Akhiezer et al., Uspekhi fiz. nauk, 71, 533 (1960)). In the approximation in question (second approximation in perturbation theory), strong exchange interaction does not contribute to absorption. It can be seen from the (1) and (2) that the transition matrix elements that are linear with respect to the variable field are nonvanishing only in second and higher perturbation-theoretical approximation. The corresponding lowest order processes are: (a) merging of a photon and a spin wave, which form two spin waves and (b) merging of two spin waves and one photon to form one spin wave. The

Card 2/4

30064 S/048/61/025/011/008/031
B108/B138

An absorption mechanism for a...

calculation of the appropriate absorption coefficients Γ_a and Γ_b is similar to that given in detail in Ref. 3 (Kaganov M. I., Tsukernik V. M., present number of this periodical, p. 1346). In this article, only the final results for the total absorption coefficient $\Gamma = \Gamma_a + \Gamma_b$ are given, with precision up to a numerical factor of the order of unity. At high temperatures ($\epsilon_0 \ll T \ll \theta_c$)

$$\Gamma \sim \begin{cases} gM_0 \frac{w^2}{\epsilon_0 \theta_c} \left(\frac{T}{\theta_c} \right)^2, & \frac{\hbar}{\tau_{ee}} \ll \hbar\omega \ll \epsilon_0, \\ gM_0 \frac{w^2}{\hbar\omega \theta_c} \left(\frac{T}{\theta_c} \right)^2, & \epsilon_0 \ll \hbar\omega \ll T, \\ gM_0 \frac{w^2}{\theta_c^{3/2} (\hbar\omega)^{1/2}} \left(\frac{T}{\theta_c} \right)^{3/2}, & T \ll \hbar\omega \ll \theta_c. \end{cases}$$

At low temperatures ($T \ll \epsilon_0$)

Card 3/4

An absorption mechanism for a...

30064 S/048/61/025/011/008/031
B108/B138

$$\Gamma \sim \begin{cases} gM_0 \frac{w^4}{\theta_c^2} \left(\frac{\epsilon_0}{\theta_c} \right)^{1/2} \left(\frac{T}{\theta_c} \right)^{1/2} e^{-\frac{3\epsilon_0}{T}}, & \frac{\hbar}{\tau_{11}} \ll \hbar\omega \ll \epsilon_0, \\ gM_0 \frac{w^2}{\theta_c^{1/2} (\hbar\omega)^{1/2}} \left(\frac{T}{\theta_c} \right)^{1/2} e^{-\frac{\epsilon_0}{T}}, & \hbar\omega \gg \epsilon_0. \end{cases}$$

Comparison of the results obtained with the absorption coefficient due to the disintegration of one photon into two spin waves (Ref. 1) shows that the processes considered in this paper are only effective at $\hbar\omega < 2\epsilon_0$.
[Abstracter's note: Complete translation.] This paper was read at the Conference on ferromagnetism and antiferromagnetism in Leningrad, May 5-11, 1961. There are 3 Soviet references.

ASSOCIATION: Fiziko-tekhnicheskiy institut Akademii nauk USSR
(Physicotechnical Institute of the Academy of Sciences
UkrSSR)

Card 4/4

24.2200

26)24.
S/056/61/041/001/019/021
B102/B231

AUTHORS: Kaganov, M. I., Tsukernik, V. M.

TITLE: High-frequency magnetic susceptibility of uniaxial ferromagnetic crystal in a longitudinal magnetic field

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 41, no. 1(7), 1961, 267 - 271

TEXT: Susceptibility dispersion of an antiferromagnetic material is mainly due to rotation of the magnetic moments of the sublattices in a magnetic field. The imaginary part of χ exhibits a resonance character in this case: resonance frequency coincides with the natural rotation frequencies of the system of moments. These frequencies have been computed for various equilibrium configurations by C. Kittel (Phys. Rev., 85, 329, 1952) and Ye. A. Turov (ZhETF, 34, 1009, 1958). In an earlier paper (ZhETF, 24, 524, 1958), the authors of the present work have computed the h-f susceptibility of a uniaxial antiferromagnetic body without magnetic field. If the magnetic field is strong enough, the equilibrium configuration of the sublattice moments may change, which

Card 1/7

26424

S/056/61/041/001/019/021

B102/B231

High-frequency magnetic ...

would result in a change of the dispersion character. Starting from the Landau-Lifshits equations for the motion of the sublattice moments, the h-f susceptibility tensor of an antiferromagnetic body is computed for various values of the constant magnetic field strength, that is, for various equilibrium configurations of the moments. The results permit to draw conclusions from h-f measurements as to the equilibrium structures and the transitions between them. The uniaxial antiferromagnetic body is assumed to have two sublattices and to be positioned in a homogeneous, constant and in a weak and variable (frequency ω) magnetic field. The motion of the moments due to field action is described by

$$\partial \vec{M}_s / \partial t = g [\vec{M}_s \vec{H}_e^{(s)}] - (\gamma / M_s^2) [\vec{M}_s [\vec{M}_s \vec{H}_e^{(s)}]] \quad (1),$$

where \vec{M}_s is the magnetization vector of the s-th sublattice, M the sublattice magnetization which is assumed to be constant, g the gyromagnetic ratio, γ the relaxation constant, $\vec{H}_e^{(s)}$ the effective field acting upon the s-th sublattice: $\vec{H}_e^{(s)} = -\partial \chi / \partial \vec{M}_s$, where χ stands for the energy density of the antiferromagnetic body:

Card 2/7

High-frequency magnetic ...

26424
S/056/61/041/001/019/021
B102/B231

$$\mathcal{H} = \alpha M_1 M_2 - \frac{1}{2} \lambda [(M_1 n)^2 + (M_2 n)^2] + \eta (M_1 n) (M_2 n) - H (M_1 + M_2). \quad (3)$$

Here, α is the constant of exchange interaction ($\alpha > 0$), λ and η represent the anisotropy constants which are assumed to be positive; n is the unit vector in the direction of the axis of the antiferromagnetic body. Using the notation stated hereinafter: $H_1 = \sqrt{(\lambda + \eta)(2\alpha - \lambda + \eta)} M$,

$$H_2 = \sqrt{(\lambda + \eta)(2\alpha + \lambda + \eta)} M, \quad H = H_3 = (2\alpha - \lambda + \eta) M, \text{ and}$$

$$H_4 = [\lambda(2\alpha - \lambda + \eta)^2 / (2\alpha + \lambda + \eta)]^{1/2} M \quad (H_4 < H_1; H_2 - H_4 \text{ is the width of}$$

the hysteresis loop of the antiferromagnetic body), the following cases are subjected to investigation: 1) $H < H_1$. m_{\pm} is equal to $\chi_{\pm} h_{\pm}$, where

$h_{\pm} = h_x \pm i h_y$, $m_{\pm} = m_x \pm i m_y$, \vec{h} stands for the h-f magnetic field, and \vec{m} for the variable part of the entire magnetic moment. With

$$\Omega^2 = (g^2 M^2 + \gamma^2) (\lambda + \eta) (2\alpha + \lambda + \eta) - \gamma^2 H^2 / M^2, \quad (5)$$

$$\Omega_1^2 = 2 (g^2 M^2 + \gamma^2) (\lambda + \eta).$$

Card 3/7

26424
S/056/61/041/001/019/021
B102/B231

High-frequency magnetic ...

the following is obtained:

$$\chi_{\pm} = \frac{\Omega_1^2 - 2i\omega\gamma}{\Omega^2 - (\omega \mp gH)^2 - 2i\omega\gamma(\alpha + \lambda + \eta)}, \quad \chi_{zz} = 0. \quad (6)$$

The width of the antiferromagnetic resonance lines is thus given by $2\gamma(\alpha + \lambda + \eta)$. With $H = 0$ one has $\chi_+ = \chi_- = \chi_{xx} = \chi_{yy}$. 2) $H_1 < H < H_3$;

Here, $\chi_{zz} \neq 0$, and the following is obtained:

$$\chi_{zz}(\omega) = \chi_{zz}(0) \frac{\nu^2 + i\nu\omega}{\nu^2 + \omega^2}; \quad (7)$$

$$\chi_{zz}(0) = 2/(2\alpha - \lambda + \eta). \quad (8)$$

$$\nu = (2\alpha - \lambda + \eta) \gamma \sin^2 \theta = \gamma (1 - H^2/H_3^2) H_3/M.$$

From this, the following is obtained for the behavior of the relaxation time $\tau = 1/\nu$ in the vicinity of the point of second-kind phase transition (with $H \approx H_3$):

$$\tau = \frac{1}{\gamma} \frac{H_3}{H_3^2 - H^2}. \quad \text{If } T \text{ is fixed, and } H = H_3 \text{ the following is valid:}$$

Card 4/7

26424
S/056/61/041/001/019/021
B102/B231

High-frequency magnetic ...

$$\tau = \frac{G}{|T_K - T|}, \quad G = M/2\gamma \left| \frac{dH_s}{dF} \right|_{T=T_K} \quad (8''),$$

where T_K is the temperature of phase transition. This means that with the frequency being fixed there is a field or a temperature at which the h-f field (polarized along z) exhibits an absorption maximum whose position and height is dependent on ω . The transverse components are in this case given by

$$\begin{aligned} \chi_{xx} &= \frac{\omega_1^2 - 2i\omega\gamma}{\omega_0^2 - \omega^2 - 2i\alpha\gamma'\omega} \cos^2 \theta, \\ \chi_{yy} &= \frac{1}{\alpha} \frac{\omega_0^2 - 2i\omega\gamma}{\omega_0^2 - \omega^2 - 2i\alpha\gamma'\omega}, \\ \chi_{xy} &= -\chi_{yx} = \frac{2igM\omega \cos \theta}{\omega_0^2 - \omega^2 - 2i\alpha\gamma'\omega}. \end{aligned} \quad (9),$$

Card 5/7

26: 24
S/056/61/041/001/019/021
B102/B231

High-frequency magnetic ...

where

$$\begin{aligned}\omega_0^2 &= (g^2 M^2 + \gamma^2) (4\alpha^2 \cos^2 \theta - 2(\lambda + \eta) \alpha \sin^2 \theta), \\ \omega_1^2 &= 4\alpha (g^2 M^2 + \gamma^2), \quad \gamma' = \gamma [1 + \cos^2 \theta + \frac{1}{2\alpha} (\lambda + \eta) \sin^2 \theta].\end{aligned}\quad (10)$$

3) $H > H_3$, $(\vec{H} \parallel \vec{n})$. Here, the tensor of h-f susceptibility coincides with that of the susceptibility of the uniaxial antiferromagnetic body:

$$\begin{aligned}\chi_{xx}(\omega) &= \chi_{yy}(\omega) = \chi_L(0) \frac{\omega_\phi^2 - i\omega\gamma_\phi}{\omega_\phi^2 - \omega^2 - 2i\omega\gamma_\phi}, \\ \chi_{xy}(\omega) &= -\chi_{yx}(\omega) = \frac{2igM\omega}{\omega_\phi^2 - \omega^2 - 2i\omega\gamma_\phi},\end{aligned}\quad (11),$$

$$\chi_{xz} = \chi_{yz} = \chi_{zx} = 0,$$

$$\begin{aligned}\chi_L(0) &= 2M/(H + (\lambda - \eta)M), \\ \omega_\phi^2 &= g^2 (H + (\lambda - \eta)M)^2 (1 + \gamma^2/g^2 M^2), \\ \gamma_\phi &= \gamma (H/M + \lambda - \eta).\end{aligned}\quad (12).$$

Card 6/7

High-frequency magnetic ...

26424
S/056/61/041/001/019/021
B102/B231

With $H = H_3 = (2\alpha - \lambda + \eta)M$ all components of $\chi_{1k}(\omega)$ are continuous.
The authors thank A. S. Borovik-Romanov for remarks. There are
7 references: 3 Soviet-bloc and 4 non-Soviet-bloc.

ASSOCIATION: Fiziko-tehnicheskii institut Akademii Nauk Ukrainskoy SSR
(Institute of Physics and Technology of the Academy of
Sciences Ukrainskaya SSR)

SUBMITTED: February 17, 1961

Card 7/7

24.4/00

S/056/62/043/003/037/063
B108/B102

AUTHORS: Gurzhi, R. N., Tsukernik, V. M.

TITLE: High-frequency magnetic susceptibility of ferromagnetic dielectrics in a longitudinal magnetic field

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, . . .
v. 43, no. 3(9), 1962, 977 - 983

TEXT: Using a method-similar to Bogolyubov's (ZhETF, 17, 614, 1947) the authors got the following quantum kinetic equation (linear approximation with regard to the variable field $h_t = h_0 e^{-i\omega t}$) for the distribution function $F_k = \langle a_k^+ a_k \rangle$ of spin waves:

$$-i\omega f_k = L_k h_0 + \hat{W} f_k; \quad (11)$$

$$L_k = (\pi\mu\omega/2) |B_k|^2 (2\tilde{n}_k + 1) \tilde{e}_k^{-2} [\delta_- (-2\tilde{e}_k - \hbar\omega) - \delta_- (2\tilde{e}_k - \hbar\omega)] + \\ + 4\pi\mu\omega \sum_{\mathbf{l}} \frac{|\Phi_{k\mathbf{l}; 2}|^2}{e_{k\mathbf{l}; 2}^2} [(1 + n_k + n_{\mathbf{l}}) n_2 - n_{\mathbf{l}} n_k] [\delta_- (e_{2; \mathbf{l}k} - \hbar\omega) - \delta_- (e_{k\mathbf{l}; 2} - \hbar\omega)] +$$

Card 1/4

High-frequency magnetic...

S/056/62/043/003/037/063
B108/B102

$$\begin{aligned}
 & + 2\pi i \omega \sum_{12} \frac{|\Phi_{12; k}|^2}{e_{12; k}^2} [(1 + n_1 + n_2)n_k - n_1 n_2] [\delta_-(e_{12; k} - \hbar\omega) - \delta_-(e_{k; 12} - \hbar\omega)], \\
 \hat{W}f_k = & \frac{4\pi}{\hbar} \sum_{12} |\Phi_{k1; 2}|^2 [(1 + n_k + n_1)f_2 + (n_2 - n_k)f_1 + (n_2 - n_1)f_k] \times \\
 & \times [\delta_-(e_{k1; 2} - \hbar\omega) + \delta_-(e_{2; k1} - \hbar\omega)] + \frac{2\pi}{\hbar} \sum_{12} |\Phi_{12; k}|^2 [(n_2 - n_k)f_1 + \\
 & + (n_1 - n_k)f_2 - (1 + n_1 + n_2)f_k] [\delta_-(e_{k; 12} - \hbar\omega) + \delta_-(e_{12; k} - \hbar\omega)] + \\
 & + \frac{16\pi}{\hbar} \sum_{123} |\Phi_{k1; 23}|^2 \{ [n_1(n_2 + 1)(n_3 + 1) - (n_1 + 1)n_2 n_3] f_k + [n_k(n_2 + 1) \times \\
 & \times (n_3 + 1) - (n_k + 1)n_2 n_3] f_1 + [n_k n_1(n_2 + 1) - (n_k + 1)(n_1 + 1)n_2] f_2 + \\
 & + [n_k n_1(n_2 + 1) - (n_k + 1)(n_1 + 1)n_2] f_3 \} [\delta_-(e_{k1; 23} - \hbar\omega) + \\
 & + \delta_-(e_{23; k1} - \hbar\omega)].
 \end{aligned}$$

Card 2/4

High-frequency magnetic...

S/056/62/043/003/037/063
B108/B102

with $\bar{r}_\alpha|_{t=0} = n_\alpha = (e^{\epsilon_\alpha/T} - 1)^{-1}$ and $\langle a_\alpha^\dagger a_\beta^\dagger a_\gamma \rangle|_{t=0} = 0$.

$\epsilon_k = \theta_{\text{Curie}}(ak)^2 + \mu H + |B_k|$. At high frequencies it is sufficient to consider the term $-i\omega f_k = L_k h_0$. From this equation the high-frequency magnetic susceptibility of a ferroelectric with one magnetic sublattice can be calculated keeping in mind that magnetization too can be expressed in terms of the spin wave distribution function:

$$V^{-1} \langle M_z \rangle = M_0 - \frac{\mu}{V} \sum_k \langle a_k^\dagger a_k \rangle, \quad \langle a_k^\dagger a_k \rangle = \text{Sp } \rho a_k^\dagger a_k, \quad (6).$$

Thus, $\chi = -\frac{i\mu}{\omega V} \sum_k L_k$. V is the volume of the body. The real part of magnetic susceptibility which is related to dispersion is

$$\text{Re } \chi = \frac{\mu}{\omega V} \sum_k \text{Im } L_k.$$

The predominant term in $\text{Im } L_k$ is the one proportional to $|B_k|^2$. Con-

Card 3/4

High-frequency magnetic...

S/056/62/043/003/037/063
B108/B102

sidering this and using the symbols $x = \sqrt{\theta_{\text{Curie}}/T} ak$, $\alpha = \hbar\omega/T$,
 $\eta = \mu H/T$, $\gamma = 4\pi\mu M_0/T$ and $w = \mu^2/a^3$ the result is

$$\text{Re } \chi = \frac{w}{4T} \left(\frac{\mu M_0}{T} \right)^2 \left(\frac{T}{\theta_c} \right)^{1/2} \int_0^\pi \sin^2 \Theta d\Theta \int_{-\infty}^{\infty} \frac{\text{ch}(y(x)/2)}{y(x)} \frac{x^2 dx}{y^2(x) - a^2/4}$$

$$y(x) = \sqrt{(x^2 + \eta)^2 + \gamma(x^2 + \eta) \sin^2 \Theta}.$$

ASSOCIATION: Fiziko-tekhnicheskii institut Akademii nauk Ukrainsskoy SSR
(Physicotechnical Institute of the Academy of Sciences of the
Ukrainsskaya SSR)

SUBMITTED: March 31, 1962

Card 4/4

L 40173-66 EWT(1) JW

ACC NR: AP6018817 SOURCE CODE: UR/0056/66/050/005/1377/1380

AUTHOR: Pikin, S. A.; Tsukernik, V. M.

ORG: none

TITLE: Thermodynamics of linear spin chains in a transverse magnetic field

SOURCE: Zh eksper i teor fiz, v. 50, no. 5, 1966, 1377-1380

TOPIC TAGS: spin system, transverse magnetic field, magnetic susceptibility, THERMODYNAMIC PROPERTY, HOMOGENEOUS MAGNETIC FIELD

ABSTRACT: A one-dimensional system of spins ($s = 1/2$) with a strongly anisotropic interaction of the immediate neighbors has been examined. The system is assumed to be in a homogeneous magnetic field directed along a given axis. Exact equations have been obtained for the thermodynamic characteristics of the system. It has been shown that at $T = 0$ and an absolute value of the magnetic-field strength H_0 , the magnetic susceptibility has a logarithmic or root nature, dependent, like H_0 , on the relation between the interaction constants. The authors thank V. G. Vaks and A. I. Larkin for valuable discussions. Orig. art. has: 11 formulas. [Based on authors' abstract] [NT]

SUB CODE: 20/ SUBM DATE: 03Dec65/ ORIG REF: 003/ OTH REF: 002

L 41744-66 EWT(1) IJP(c) GG

ACC NR: AP6020219

SOURCE CODE: UR/0056/66/050/006/1631/1635

AUTHOR: Tsukernik, V. M.

ORG: none

TITLE: Features of gyromagnetic effect in ferroelectrics at low temperatures

SOURCE: Zh eksper i teor fiz, v. 50, no. 6, 1966, 1631-1635

TOPIC TAGS: ferromagnetic dielectric, spin wave theory, dipole interaction, magnetic moment, ground state, temperature dependence

ABSTRACT: The author derives in the spin-wave approximation a formula for the temperature part of the mechanical moment of a spin system and shows that in the region of temperatures where the dipole interaction plays an important role, its ratio to the corresponding addition to the magnetic moment differs from the single-electron g factor and depends on the temperature. The approximation employed consists of expanding, with accuracy to quadratic terms, the energy density in powers of a quantity which is small near the ground state in comparison with the magnetization, followed by quantization and averaging with an equilibrium Gibbs distribution. The deduced difference in the temperature dependence of the magnetic and mechanical moments of the spin system is observable in principle with the aid of a gyromagnetic experiment in which the moments are changed as a result of changes in the temperature in the absence of an external field. The author thanks M. I. Kaganov for useful discussions. Orig. art. has: 20 formulas.

SUB CODE: 20/ SUBM DATE: 18Jan65/ ORIG REF: 004

Card 1/1

SOURCE: Zhurnal eksperimental'noy i teoreticheskoy fiziki,
v. 48, no. 5, 1965, 1386-1392

TOPIC TAGS: electromagnetic field, field state, field intensity
value, wave function, occupation number, quantum electrodynamics

ABSTRACT: Since a free electromagnetic field can be described in
quantum theory either by giving the photon occupation numbers or
by giving the field strength, the authors consider the problem of
finding states of the electromagnetic field in which the electrical
or the magnetic field strength has a well defined value. The states
of interest are those in which the field strength is given in all of
space at one particular time, so that a complete set of the values
of the electrical (or magnetic) field strength can be chosen in all

skoy SSR (Institute of Radiophysics and Electronics, Academy of Sciences of the Ukrainian SSR)

NR REF SOV: 003

OTHER: 002

ZAKHAROV, A.G.; VECHER, N.A.; LEKONISEV, A.N.; RUDNITSKIY, P.M.;
TOLKALINIKO, L.N.; TSHEPNIK, Z.G.; ARYASOV, N.I., inzh.,
inzhent; DOVCOFOL, V.I., red.; DEBROV, I.P., red.;
GETLING, Yu., red.

[Vanadium of the Kachkanar deposit] Kachkanarskii vanadii.
Sverdlovsk, Sredne-Ural'skoe knizhnoe izd-vo, 1964. 302 p.
(MIRA 18:11)

BURDAKOV, Dmitriy Dmitriyevich; TSUKERNIK, Zinoviy Grigor'yevich; YEL'KIN, S.A., inzh., retsenzent; ROMANOV, A.A., kand. tekhn. nauk, retsenzent; BENYAKOVSKIY, M.A., inzh., retsenzent; GOL'DSHTEYN, M.I., kand. tekhn. nauk, retsenzent; DUBROV, N.F., nauchnyy red.; SYRCHINA, M.M., red. izd-va; KRYZHOVA, M.L., red. izd-va; TURKINA, Ye.D., tekhn. red.

[Metallurgy of ferrous metals; manual for the training of skilled workers in industry] Metallurgiya chernykh metallov; uchebnoe posobie dlia podgotovki kvalifitsirovannykh rabochikh na proizvodstv. Sverdlovsk, Gos. nauchno-tekhn. izd-vo lit-ry po chernoi i tsvetnoi metallurgii, 1961. 437 p. (MIRA 14:11)
(Iron—Metallurgy) (Steel—Metallurgy) (Metalwork)

IGNATIK, A.F., inzh.; TSUKERNIK, Z.S., inzh.

Using radioactive pickups in the automatization and regulation
of stone-materials production. Stroi.i dor.mashinostr. 4 no.9:
28-30 S '59. (MIRA 12:11)

(Radioisotopes--Industrial applications)

(Automatic control)

(Building materials industry)

IGNATYUK, A.P., inzh.; TSUKERNIK, Z.S., inzh.

Using tagged atoms for determining the speed of movement of
concrete through steel pipes. Stroil. i dor. mashinostr 3 no.5:
23-24 My '58. (MIRA 11:5)
(Radioactive tracers--Industrial applications)
(Concrete--Transportation)

— TSUKERNIK, Z.S.

TSUKERNIK, Z.S., inzhener.

New tensiometering instrument. Sbor.trud.VNIISTroidorrash
Lenfil. no.1:91-95 '57. (PIRA 10:8)
(Electronic instruments)
(Tensiometers)

PANFILOVA, T.A.; SUVOROVSKAIA, N.A.; TSUKERSHTEYN, O.E.

The coronary vasoconstrictor activity of the serum of rabbits
on a cholesterol diet. Cor Vasa 6 no.4:308-311 '64.

1. The Central Research Laboratory, 1st Leningrad Medical
Institute, Leningrad, USSR.

PANFILOVA, T.A.; TSUKERSHTEYN, O.Ya.

Modified spreading rate of the pulse wave in some arrhythmias under experimental conditions. Biul.eksp.biol.i med. 58 no.7:30-32 J1 '64.
(MIRA 18:2)

1. Tsentral'naya nauchno-issledovatel'skaya laboratoriya (zav. - dotsent V.M.Ivanov) I Leningradskogo meditsinskogo instituta imeni Pavlova. Submitted June 15, 1963.

TSUKERSHTEYN, O.Ye.

Effect of vagotropic substances on certain functions of the
liver, effect of the nervous system on the liver. Ter. arkh.,
Moskva 23 no. 6:43-55 Nov-Dec. 1951. (CIML 21:3)

1. Of the Naval Medical Academy..

TSUKERSHTEYN, O.Ye.; KUPSIK, B.M.

Some variants in the hemopoietic system in diphyllbothriasis.
Probl.gemat. i perel.krovi 4 no.7:49-52 J1 '59. (MIRA 12:10)

1. Iz I Leningradskogo meditsinskogo instituta imeni akad.
I.P.Pavlova.

(TAPEWORM INFECTION, compl.
diphyllbothriasis causing anemia (Rus))
(ANEMIA, etiol. & pathogen.
diphyllbothriasis (Rus))

TSUKERSHTEYN, O. Ye.
TSUKERSHTEYN, O. Ye.; KUPSIK, B.M.

Some peculiarities in the composition of the blood in diphyllor-
bothriasis in the Far North. Sov.med. 21 Supplement:7 '57.
(ARTIC REGIONS--~~TAP~~WORMS) (MIRA 11:2)
(BLOOD--~~EXAMINATION~~)

LINDENBRATEN, L.D. (Leningrad); TSUKERSHTEYN, O.Ye. (Leningrad).

Diagnosis of internal biliary fistulas; symptom of gas in bile ducts. Klin.
med. 31 no.8:38-42 Ag '53. (MLRA 6:11)

(Biliary tract) (Fistula)

TSUKERSHTEYN, O.Ye., kand.med.nauk

Case of diabetes mellitus developing under the influence of ACTH
and cortisone. Probl.endok.i gorm. no.1:115-116 '62. (MIRA 15:8)

(ACTH)

(CORTISONE)

(DIABETES)

PANFILOVA, T.A.; TSUKERSHTEYN, O.Ye.; VOL'FE, M.M.

Correlations between functional and morphological changes in
experimental cholesterol atherosclerosis in dogs. Pat. fiziol.
i eksp. terap. 9 no.5:53-57 S-O '65. (MIRA 19:1)

1. "Sentral'naya nauka i issledovatel'skaya laboratoriya (zav. -
dozent V.M. Ivanov) i Leningradskogo meditsinskogo instituta
imeni I.P. Pavlova. Submitted May 23, 1964.

TSUKERSHTEYN, O. YE.

"Leukemoid Reactions During Intoxication With Tetraethyl Lead",
Military-Medical Journal, No. 8, P 34, Aug 1955.

MASLENNIKOV, V.D.; TSUKERSHTEYN, O.Ye.

Leucocytosis and absolute monocytosis following Valdman's cup
endothelial test. Lab. delo 7 no.3:22-25 Mr '61. (MIRA 14:3)

(LEUCOCYTOSIS)

(RHEUMATIC FEVER)

TSUKERSHTEYN, V.G., and RUDERMAN, A.I.

"Experimental Studies on the Distribution of X-ray Radiation Doses when Sectorial Irradiation by Means of Swinging X-ray Tubes is being applied", p. 165. Research Into the Field of X-Ray Technique, Vol. 9, Medgiz, Moscow 1955

RUDERMAN, A.I., doktor med. nauk; TSUKERSHTEYN, V.G..

Indications for X-ray treatment of esophageal cancer by sector
irradiation with a pivoting tube. Trudy TSentr. nauch.-issl. inst.
rentg. 1 rad. 10:226-230 '59. (MIRA 12:9)
(ESOPHAGUS--CANCER) (X RAYS--THERAPEUTIC USE)

TSUKERSHTEYN, V.G.

X-ray treatment of benign vascular tumors. Trudy TSentr. nauch.-
issl. inst. rentg. 1 rad. 10:273-278 '59. (MIRA 10:9)
(X RAYS---THERAPEUTIC USE) (TUMORS)

RUDERMAN, A.I.: TSUKERSHTEYN, V.G.

Experimental study of the distribution of X-ray emission in the case
of sectional exposure by oscillating tubes. Trudy TSentr. nauchn.
issl.inst.rentg. 1 rad. 9:165-173 '55. (MLBA 9:12)
(X RAYS)

TSUKERSHTEIN, ⁴E. I.

Diabetes 2. izd-perer. i dop. Leningrad, Gos. ordena Lenina in-t
usovershenstvovaniia vrachei im. S.M. Kirova, 1947. 336 P.

DAFM

L. Diabetes

PA 65/49168

TSUKERSHTEYN, Ye.I. (Prof)

USSR/Medicine - Blood, Cholesterol * Mar 49
Blood, Ketone Bodies

"Presence of Ketone Bodies and Cholesterol in the Blood of Hypertonia Cases," Prof Ye. I. Tsukershteyn, Chief, Third Chair of Internal Diseases, M. N. Yegorova, Third Chair of Internal Diseases, State Ord of Lenin Inst for Advancement of Doctors (men) S. M. Kirov, 6 pp

"Klin Med" No 3

Gives results of studying hypertonic cases, in three tables. From data obtained, it may be

65/49168

USSR/Medicine - Blood, Cholesterol Mar 49
(Contd.)

assumed that hyperketonemia has a serious effect on the general condition of the patient. Changes in general or alimentary functions affect the content of ketone bodies. In most cases under proper treatment, the cholesterol content in the blood tends to approach normal as general condition and liver functions improve. Dir, State Ord of Lenin Inst for Advancement of Doctors: G. A. Znamenskiy.

65/49168

TSUKERSHTEYN, Ye.I.; YNGOROVA, M.N.

Barrier liver function in hypertension. Klin. med., Moskva 30 no.
11:90 Nov 1952. (GLML 23:5)

1. Professor for Tsukershteyn. 2. Of the Third Department of Internal Diseases (Head -- Prof. Ye. I. Tsukershteyn), State Order of Lenin Institute for the Advanced Training of Physicians imeni S. M. Kirov.

TSUKERSHTEYN, Ye.I., professor (Leningrad)

Peculiarities of the course of rheumatism in adolescents. Klin. med.
31 no.11:48-52 N '53.

(MLRA 6:12)
(Rheumatism)

TSUKERSHTEYN, Ye.I., prof.; TNIMOVA, K.Z., dotsent

Peculiarities in the clinical aspects and treatment of tuberculosis of the lymph nodes of the abdominal cavity. Zdrav. Kazakh. 21 no.1: 43-48 '61. (MIRA 14:3)

1. Iz kafedry gospi'tal'noy terapii (zav. prof. Ye.I. Tsukershteyn)
Karagandinskogo meditsinskogo instituta.
(LYMPHATICS—TUBERCULOSIS) (ABDOMEN)

TSUKERSHTEYN, Ye.I., prof.; KARLINSKIY, V.M., kand.med.nauk

Prevention of rheumatic fever in adults. *Zdrav. Kazakh*, 21 no.5:
5-8 '61. (MIRA 15:2)

1. Iz kafedry gosptal'noy terapii Karagandinskogo meditsinskogo
instituta.

(RHEUMATIC FEVER)

A-3

117 APR 199 200131


PROCESSES AND PROPERTIES INDEX

BE

Characterization of quinols. I. TSURUMIYANER
(Bull. Univ. Asia Central, 1934, No. 18, 48-50).
Tolquinol and Br in CHCl_3 yield the dibromide, an
oil, readily decomp. to yield 2-chloro-5-methylquinol;
the tetrabromide, m.p. 118°, prepared similarly, yields
di- and tri-bromo-p-quinol when heated above the
m.p., whilst when (2) is heated with H_2O the product
is 2:6-dibromo-5-methylquinol. The di-, m.p. 93°
(decomp.), and tetrabromide, m.p. 145°, prepared
analogously, behave similarly to the above bromides,
but are more stable. Epoxidized dibromide, m.p.
108°, tetrabromide, m.p. 145°, dibromide, m.p. 112°,
and tetrachloride, an oil, yield unidentified quinols
when heated: (3) (4) (5) (6) (7) (8) (9) (10) (11) (12) (13) (14) (15) (16) (17) (18) (19) (20) (21) (22) (23) (24) (25) (26) (27) (28) (29) (30) (31) (32) (33) (34) (35) (36) (37) (38) (39) (40) (41) (42) (43) (44) (45) (46) (47) (48) (49) (50) (51) (52) (53) (54) (55) (56) (57) (58) (59) (60) (61) (62) (63) (64) (65) (66) (67) (68) (69) (70) (71) (72) (73) (74) (75) (76) (77) (78) (79) (80) (81) (82) (83) (84) (85) (86) (87) (88) (89) (90) (91) (92) (93) (94) (95) (96) (97) (98) (99) (100) (101) (102) (103) (104) (105) (106) (107) (108) (109) (110) (111) (112) (113) (114) (115) (116) (117) (118) (119) (120) (121) (122) (123) (124) (125) (126) (127) (128) (129) (130) (131) (132) (133) (134) (135) (136) (137) (138) (139) (140) (141) (142) (143) (144) (145) (146) (147) (148) (149) (150) (151) (152) (153) (154) (155) (156) (157) (158) (159) (160) (161) (162) (163) (164) (165) (166) (167) (168) (169) (170) (171) (172) (173) (174) (175) (176) (177) (178) (179) (180) (181) (182) (183) (184) (185) (186) (187) (188) (189) (190) (191) (192) (193) (194) (195) (196) (197) (198) (199) (200) (201) (202) (203) (204) (205) (206) (207) (208) (209) (210) (211) (212) (213) (214) (215) (216) (217) (218) (219) (220) (221) (222) (223) (224) (225) (226) (227) (228) (229) (230) (231) (232) (233) (234) (235) (236) (237) (238) (239) (240) (241) (242) (243) (244) (245) (246) (247) (248) (249) (250) (251) (252) (253) (254) (255) (256) (257) (258) (259) (260) (261) (262) (263) (264) (265) (266) (267) (268) (269) (270) (271) (272) (273) (274) (275) (276) (277) (278) (279) (280) (281) (282) (283) (284) (285) (286) (287) (288) (289) (290) (291) (292) (293) (294) (295) (296) (297) (298) (299) (300) (301) (302) (303) (304) (305) (306) (307) (308) (309) (310) (311) (312) (313) (314) (315) (316) (317) (318) (319) (320) (321) (322) (323) (324) (325) (326) (327) (328) (329) (330) (331) (332) (333) (334) (335) (336) (337) (338) (339) (340) (341) (342) (343) (344) (345) (346) (347) (348) (349) (350) (351) (352) (353) (354) (355) (356) (357) (358) (359) (360) (361) (362) (363) (364) (365) (366) (367) (368) (369) (370) (371) (372) (373) (374) (375) (376) (377) (378) (379) (380) (381) (382) (383) (384) (385) (386) (387) (388) (389) (390) (391) (392) (393) (394) (395) (396) (397) (398) (399) (400) (401) (402) (403) (404) (405) (406) (407) (408) (409) (410) (411) (412) (413) (414) (415) (416) (417) (418) (419) (420) (421) (422) (423) (424) (425) (426) (427) (428) (429) (430) (431) (432) (433) (434) (435) (436) (437) (438) (439) (440) (441) (442) (443) (444) (445) (446) (447) (448) (449) (450) (451) (452) (453) (454) (455) (456) (457) (458) (459) (460) (461) (462) (463) (464) (465) (466) (467) (468) (469) (470) (471) (472) (473) (474) (475) (476) (477) (478) (479) (480) (481) (482) (483) (484) (485) (486) (487) (488) (489) (490) (491) (492) (493) (494) (495) (496) (497) (498) (499) (500) (501) (502) (503) (504) (505) (506) (507) (508) (509) (510) (511) (512) (513) (514) (515) (516) (517) (518) (519) (520) (521) (522) (523) (524) (525) (526) (527) (528) (529) (530) (531) (532) (533) (534) (535) (536) (537) (538) (539) (540) (541) (542) (543) (544) (545) (546) (547) (548) (549) (550) (551) (552) (553) (554) (555) (556) (557) (558) (559) (560) (561) (562) (563) (564) (565) (566) (567) (568) (569) (570) (571) (572) (573) (574) (575) (576) (577) (578) (579) (580) (581) (582) (583) (584) (585) (586) (587) (588) (589) (590) (591) (592) (593) (594) (595) (596) (597) (598) (599) (600) (601) (602) (603) (604) (605) (606) (607) (608) (609) (610) (611) (612) (613) (614) (615) (616) (617) (618) (619) (620) (621) (622) (623) (624) (625) (626) (627) (628) (629) (630) (631) (632) (633) (634) (635) (636) (637) (638) (639) (640) (641) (642) (643) (644) (645) (646) (647) (648) (649) (650) (651) (652) (653) (654) (655) (656) (657) (658) (659) (660) (661) (662) (663) (664) (665) (666) (667) (668) (669) (670) (671) (672) (673) (674) (675) (676) (677) (678) (679) (680) (681) (682) (683) (684) (685) (686) (687) (688) (689) (690) (691) (692) (693) (694) (695) (696) (697) (698) (699) (700) (701) (702) (703) (704) (705) (706) (707) (708) (709) (710) (711) (712) (713) (714) (715) (716) (717) (718) (719) (720) (721) (722) (723) (724) (725) (726) (727) (728) (729) (730) (731) (732) (733) (734) (735) (736) (737) (738) (739) (740) (741) (742) (743) (744) (745) (746) (747) (748) (749) (750) (751) (752) (753) (754) (755) (756) (757) (758) (759) (760) (761) (762) (763) (764) (765) (766) (767) (768) (769) (770) (771) (772) (773) (774) (775) (776) (777) (778) (779) (780) (781) (782) (783) (784) (785) (786) (787) (788) (789) (790) (791) (792) (793) (794) (795) (796) (797) (798) (799) (800) (801) (802) (803) (804) (805) (806) (807) (808) (809) (810) (811) (812) (813) (814) (815) (816) (817) (818) (819) (820) (821) (822) (823) (824) (825) (826) (827) (828) (829) (830) (831) (832) (833) (834) (835) (836) (837) (838) (839) (840) (841) (842) (843) (844) (845) (846) (847) (848) (849) (850) (851) (852) (853) (854) (855) (856) (857) (858) (859) (860) (861) (862) (863) (864) (865) (866) (867) (868) (869) (870) (871) (872) (873) (874) (875) (876) (877) (878) (879) (880) (881) (882) (883) (884) (885) (886) (887) (888) (889) (890) (891) (892) (893) (894) (895) (896) (897) (898) (899) (900) (901) (902) (903) (904) (905) (906) (907) (908) (909) (910) (911) (912) (913) (914) (915) (916) (917) (918) (919) (920) (921) (922) (923) (924) (925) (926) (927) (928) (929) (930) (931) (932) (933) (934) (935) (936) (937) (938) (939) (940) (941) (942) (943) (944) (945) (946) (947) (948) (949) (950) (951) (952) (953) (954) (955) (956) (957) (958) (959) (960) (961) (962) (963) (964) (965) (966) (967) (968) (969) (970) (971) (972) (973) (974) (975) (976) (977) (978) (979) (980) (981) (982) (983) (984) (985) (986) (987) (988) (989) (990) (991) (992) (993) (994) (995) (996) (997) (998) (999) (1000) (1001) (1002) (1003) (1004) (1005) (1006) (1007) (1008) (1009) (1010) (1011) (1012) (1013) (1014) (1015) (1016) (1017) (1018) (1019) (1020) (1021) (1022) (1023) (1024) (1025) (1026) (1027) (1028) (1029) (1030) (1031) (1032) (1033) (1034) (1035) (1036) (1037) (1038) (1039) (1040) (1041) (1042) (1043) (1044) (1045) (1046) (1047) (1048) (1049) (1050) (1051) (1052) (1053) (1054) (1055) (1056) (1057) (1058) (1059) (1060) (1061) (1062) (1063) (1064) (1065) (1066) (1067) (1068) (1069) (1070) (1071) (1072) (1073) (1074) (1075) (1076) (1077) (1078) (1079) (1080) (1081) (1082) (1083) (1084) (1085) (1086) (1087) (1088) (1089) (1090) (1091) (1092) (1093) (1094) (1095) (1096) (1097) (1098) (1099) (1100) (1101) (1102) (1103) (1104) (1105) (1106) (1107) (1108) (1109) (1110) (1111) (1112) (1113) (1114) (1115) (1116) (1117) (1118) (1119) (1120) (1121) (1122) (1123) (1124) (1125) (1126) (1127) (1128) (1129) (1130) (1131) (1132) (1133) (1134) (1135) (1136) (1137) (1138) (1139) (1140) (1141) (1142) (1143) (1144) (1145) (1146) (1147) (1148) (1149) (1150) (1151) (1152) (1153) (1154) (1155) (1156) (1157) (1158) (1159) (1160) (1161) (1162) (1163) (1164) (1165) (1166) (1167) (1168) (1169) (1170) (1171) (1172) (1173) (1174) (1175) (1176) (1177) (1178) (1179) (1180) (1181) (1182) (1183) (1184) (1185) (1186) (1187) (1188) (1189) (1190) (1191) (1192) (1193) (1194) (1195) (1196) (1197) (1198) (1199) (1200) (1201) (1202) (1203) (1204) (1205) (1206) (1207) (1208) (1209) (1210) (1211) (1212) (1213) (1214) (1215) (1216) (1217) (1218) (1219) (1220) (1221) (1222) (1223) (1224) (1225) (1226) (1227) (1228) (1229) (1230) (1231) (1232) (1233) (1234) (1235) (1236) (1237) (1238) (1239) (1240) (1241) (1242) (1243) (1244) (1245) (1246) (1247) (1248) (1249) (1250) (1251) (1252) (1253) (1254) (1255) (1256) (1257) (1258) (1259) (1260) (1261) (1262) (1263) (1264) (1265) (1266) (1267) (1268) (1269) (1270) (1271) (1272) (1273) (1274) (1275) (1276) (1277) (1278) (1279) (1280) (1281) (1282) (1283) (1284) (1285) (1286) (1287) (1288) (1289) (1290) (1291) (1292) (1293) (1294) (1295) (1296) (1297) (1298) (1299) (1300) (1301) (1302) (1303) (1304) (1305) (1306) (1307) (1308) (1309) (1310) (1311) (1312) (1313) (1314) (1315) (1316) (1317) (1318) (1319) (1320) (1321) (1322) (1323) (1324) (1325) (1326) (1327) (1328) (1329) (1330) (1331) (1332) (1333) (1334) (1335) (1336) (1337) (1338) (1339) (1340) (1341) (1342) (1343) (1344) (1345) (1346) (1347) (1348) (1349) (1350) (1351) (1352) (1353) (1354) (1355) (1356) (1357) (1358) (1359) (1360) (1361) (1362) (1363) (1364) (1365) (1366) (1367) (1368) (1369) (1370) (1371) (1372) (1373) (1374) (1375) (1376) (1377) (1378) (1379) (1380) (1381) (1382) (1383) (1384) (1385) (1386) (1387) (1388) (1389) (1390) (1391) (1392) (1393) (1394) (1395) (1396) (1397) (1398) (1399) (1400) (1401) (1402) (1403) (1404) (1405) (1406) (1407) (1408) (1409) (1410) (1411) (1412) (1413) (1414) (1415) (1416) (1417) (1418) (1419) (1420) (1421) (1422) (1423) (1424) (1425) (1426) (1427) (1428) (1429) (1430) (1431) (1432) (1433) (1434) (1435) (1436) (1437) (1438) (1439) (1440) (1441) (1442) (1443) (1444) (1445) (1446) (1447) (1448) (1449) (1450) (1451) (1452) (1453) (1454) (1455) (1456) (1457) (1458) (1459) (1460) (1461) (1462) (1463) (1464) (1465) (1466) (1467) (1468) (1469) (1470) (1471) (1472) (1473) (1474) (1475) (1476) (1477) (1478) (1479) (1480) (1481) (1482) (1483) (1484) (1485) (1486) (1487) (1488) (1489) (1490) (1491) (1492) (1493) (1494) (1495) (1496) (1497) (1498) (1499) (1500) (1501) (1502) (1503) (1504) (1505) (1506) (1507) (1508) (1509) (1510) (1511) (1512) (1513) (1514) (1515) (1516) (1517) (1518) (1519) (1520) (1521) (1522) (1523) (1524) (1525) (1526) (1527) (1528) (1529) (1530) (1531) (1532) (1533) (1534) (1535) (1536) (1537) (1538) (1539) (1540) (1541) (1542) (1543) (1544) (1545) (1546) (1547) (1548) (1549) (1550) (1551) (1552) (1553) (1554) (1555) (1556) (1557) (1558) (1559) (1560) (1561) (1562) (1563) (1564) (1565) (1566) (1567) (1568) (1569) (1570) (1571) (1572) (1573) (1574) (1575) (1576) (1577) (1578) (1579) (1580) (1581) (1582) (1583) (1584) (1585) (1586) (1587) (1588) (1589) (1590) (1591) (1592) (1593) (1594) (1595) (1596) (1597) (1598) (1599) (1600) (1601) (1602) (1603) (1604) (1605) (1606) (1607) (1608) (1609) (1610) (1611) (1612) (1613) (1614) (1615) (1616) (1617) (1618) (1619) (1620) (1621) (1622) (1623) (1624) (1625) (1626) (1627) (1628) (1629) (1630) (1631) (1632) (1633) (1634) (1635) (1636) (1637) (1638) (1639) (1640) (1641) (1642) (1643) (1644) (1645) (1646) (1647) (1648) (1649) (1650) (1651) (1652) (1653) (1654) (1655) (1656) (1657) (1658) (1659) (1660) (1661) (1662) (1663) (1664) (1665) (1666) (1667) (1668) (1669) (1670) (1671) (1672) (1673) (1674) (1675) (1676) (1677) (1678) (1679) (1680) (1681) (1682) (1683) (1684) (1685) (1686) (1687) (1688) (1689) (1690) (1691) (1692) (1693) (1694) (1695) (1696) (1697) (1698) (1699) (1700) (1701) (1702) (1703) (1704) (1705) (1706) (1707) (1708) (1709) (1710) (1711) (1712) (1713) (1714) (1715) (1716) (1717) (1718) (1719) (1720) (1721) (1722) (1723) (1724) (1725) (1726) (1727) (1728) (1729) (1730) (1731) (1732) (1733) (1734) (1735) (1736) (1737) (1738) (1739) (1740) (1741) (1742) (1743) (1744) (1745) (1746) (1747) (1748) (1749) (1750) (1751) (1752) (1753) (1754) (1755) (1756) (1757) (1758) (1759) (1760) (1761) (1762) (1763) (1764) (1765) (1766) (1767) (1768) (1769) (1770) (1771) (1772) (1773) (1774) (1775) (1776) (1777) (1778) (1779) (1780) (1781) (1782) (1783) (1784) (1785) (1786) (1787) (1788) (1789) (1790) (1791) (1792) (1793) (1794) (1795) (1796) (1797) (1798) (1799) (1800) (1801) (1802) (1803) (1804) (1805) (1806) (1807) (1808) (1809) (1810) (1811) (1812) (1813) (1814) (1815) (1816) (1817) (1818) (1819) (1820) (1821) (1822) (1823) (1824) (1825) (1826) (1827) (1828) (1829) (1830) (1831) (1832) (1833) (1834) (1835) (1836) (1837) (1838) (1839) (1840) (1841) (1842) (1843) (1844) (1845) (1846) (1847) (1848) (1849) (1850) (1851) (1852) (1853) (1854) (1855) (1856) (1857) (1858) (1859) (1860) (1861) (1862) (1863) (1864) (1865) (1866) (1867) (1868) (1869) (1870) (1871) (1872) (1873) (1874) (1875) (1876) (1877) (1878) (1879) (1880) (1881) (1882) (1883) (1884) (1885) (1886) (1887) (1888) (1889) (1890) (1891) (1892) (1893) (1894) (1895) (1896) (1897) (1898) (1899) (1900) (1901) (1902) (1903) (1904) (1905) (1906) (1907) (1908) (1909) (1910) (1911) (1912) (1913) (1914) (1915) (1916) (1917) (1918) (1919) (1920) (1921) (1922) (1923) (1924) (1925) (1926) (1927) (1928) (1929) (1930) (1931) (1932) (1933) (1934) (1935) (1936) (1937) (1938) (1939) (1940) (1941) (1942) (1943) (1944) (1945) (1946) (1947) (1948) (1949) (1950) (1951) (1952) (1953) (1954) (1955) (1956) (1957) (1958) (1959) (1960) (1961) (1962) (1963) (1964) (1965) (1966) (1967) (1968) (1969) (1970) (1971) (1972) (1973) (1974) (1975) (1976) (1977) (1978) (1979) (1980) (1981) (1982) (1983) (1984) (1985) (1986) (1987) (1988) (1989) (1990) (1991) (1992) (1993) (1994) (1995) (1996) (1997) (1998) (1999) (2000) (2001) (2002) (2003) (2004) (2005) (2006) (2007) (2008) (2009) (2010) (2011) (2012) (2013) (2014) (2015) (2016) (2017) (2018) (2019) (2020) (2021) (2022) (2023) (2024) (2025) (2026) (2027) (2028) (2029) (2030) (2031) (2032) (2033) (2034) (2035) (2036) (2037) (2038) (2039) (2040) (2041) (2042) (2043) (2044) (2045) (2046) (2047) (2048) (2049) (2050) (2051) (2052) (2053) (2054) (2055) (2056) (2057) (2058) (2059) (2060) (2061) (2062) (2063) (2064) (2065) (2066) (2067) (2068) (2069) (2070) (2071) (2072) (2073) (2074) (2075) (2076) (2077) (2078) (2079) (2080) (2081) (2082) (2083) (2084) (2085) (2086) (2087) (2088) (2089) (2090) (2091) (2092) (2093) (2094) (2095) (2096) (2097) (2098) (2099) (2100) (2101) (2102) (2103) (2104) (2105) (2106) (2107) (2108) (2109) (2110) (2111) (2112) (2113) (2114) (2115) (2116) (2117) (2118) (2119) (2120) (2121) (2122) (2123) (2124) (2125) (2126) (2127) (2128) (2129) (2130) (2131) (2132) (2133) (2134) (2135) (2136) (2137) (2138) (2139) (2140) (2141) (2142) (2143) (2144) (2145) (2146) (2147) (2148) (2149) (2150) (2151) (2152) (2153) (2154) (2155) (2156) (2157) (2158) (2159) (2160) (2161) (2162) (2163) (2164) (2165) (2166) (21

157 AND 2ND ORDER		PROCEDURES AND PROPERTIES WORK		340 AND 4TH ORDER	
<p>BC</p>		<p>Composition of (A) <i>Psoralea drupacea</i>, L., and of the oil pressed from its seeds, (B) roots of <i>Ferula pseudocassia</i>, Lipsky. 1. TRUKH. VANIK and V. BARSUTSKI (Bull. Univ. Asie Centrale, 1933, No. 21, 49-54, 55-66).—(A) The air-dry leaves contain hemicellulose (I) 7.7, cellulose (II) 9.3, lignin (III) 5.5, protein 16.8, ash 9.7, H₂O-sol. 24, Et₂O-sol. substances 13.8%, the roots contain 8.9% of H₂O-sol. products, and the seeds contain (I) 6.8, (II) 8.3, (III) 4.1, ash 6.0, H₂O-sol. 23.5, EtOH-sol. 2.1, and Et₂O-sol. substances 15.4%. The Et₂O extract of the seeds yields a fatty oil, containing saturated (palmitic, stearic) 8, and unsaturated acids (oleic, linoleic, linolenic) 80, resins 4.1, and unsaponifiable 5.0%.</p> <p>(B) [with C. V. BRUTKINA and B. I. AJEIKOVITCH]. The air-dry root contains (I) 11, (II) 12.4, (III) 4.3, pentosans 6.4, pectins 9.6, starch 7.19, mono- 2.11, and di-saccharides 8.80, protein 7.37, ash 5.82, H₂O-sol. 29.6, EtOH-sol. 1.9, and Et₂O-sol. substances 15.9% [including resin 12.63 (acids, including podocarpic, 6.1, phenols 62.5, and neutral constituents 20.5%), essential oil 3.3 (d-α-pinene 90, d-β-phell. andrene 5%)]. R. T.</p>		<p>CL-4</p>	
<p>ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION</p>					
<p>FROM SYNDICATE</p>					
<p>FROM SYNDICATE</p>					

1ST AND 2ND ORDERS										3RD AND 4TH ORDERS									
COMMON ELEMENTS																			
COMMON VARIABLE INDEX																			
<div style="display: flex; justify-content: space-between;"> ca 13 </div> <p>Utilization of cottonseed husks for synthesis of glucose, lactic acid and glycerol. I. P. Tsukervanik, P. P. Samsonov, L. P. Danilevskii and N. G. Sidorova. <i>Bull. Acad. Sci. USSR Div. Chem. USSR</i> No. 21, 60-82(1935).—One kg. of husks is heated for 1 hr. at 20° with 1 l. of 75.5% H₂SO₄, undissolved material is removed, 8 l. of H₂O added, and the soln. is boiled (for 3 hrs.) and made neutral with CaO or CaCO₃. Glucose is obtained in 27-30% yield from the condensed filtrate and may be fermented to yield lactic acid, or EtOH and glycerol. B. C. A.</p>																			
<div style="display: flex; justify-content: space-between;"> <div> <p>ASAC-51A METALLURGICAL LITERATURE CLASSIFICATION</p> <p>REGIONAL SYMBOLISM</p> </div> <div> <p>REGIONAL SYMBOLISM</p> <p>REGIONAL SYMBOLISM</p> </div> </div>																			

1ST AND 2ND COLUMNS		PROCESS AND PROPERTIES INDEX	
<p>Alkylation of phenols with alcohols in the presence of aluminum chloride. II. Alkylation with secondary and primary aliphatic alcohols. I. P. Tsukervanik and Z. N. Nazarova. <i>J. Gen. Chem.</i> (U.S.S.R.) 7, 1237-31 (1937); <i>C. A.</i> 30, 4431. The difficulties of obtaining alkylphenols by catalytic condensation of phenols with secondary alcs. have been overcome by the use of excess AlCl_3 and higher temps. By this method about 70% alkylphenols, contg. small amts. of dialkylphenol ethers, $(\text{R}_2\text{C}_6\text{H}_3\text{OR})$ (I) and alkylphenol ethers $(\text{RC}_6\text{H}_4\text{OR})$ (II), are formed. The proportion of II in the reaction mixt. increases with decrease in the AlCl_3 concn. and temp. The probable mechanism of the condensation is: $\text{ROH} + \text{AlCl}_3 \rightarrow \text{HCl} + \text{AlCl}_2(\text{OR})$; $\text{PhOH} + \text{AlCl}_2(\text{OR}) \rightarrow \text{HCl} + \text{PhOAlCl}_2$; $\text{AlCl}_2(\text{OR}) + \text{PhOAlCl}_2 \rightarrow \text{PhOR} + \text{AlCl}_3 + \text{AlCl}_2\text{OH}$. PhOR is then converted to II either by alkylation: $\text{PhOR} + \text{ROH} + \text{AlCl}_3 \rightarrow \text{II} + \text{AlCl}_2\text{OH} + \text{HCl}$, or by rearrangement: $2 \text{PhOR} \xrightarrow{\text{AlCl}_3} \text{II} + \text{PhOH}$ (cf. Smith, <i>C. A.</i> 27, 8067). The alkylated ethers are then converted to alkylphenols by the action of the excess of AlCl_3 or higher temps. either by sapon.: $\text{II} + \text{HCl} \xrightarrow{\text{AlCl}_3} \text{RC}_6\text{H}_4\text{OH} + \text{RCI}$ (Hartmann and Gattermann, <i>Ber.</i> 25, 3331 (1892)), or by rearrangement: $\text{II} + \text{PhOH} \xrightarrow{\text{AlCl}_3}$</p>		<p>10</p> <p>$2\text{RC}_6\text{H}_4\text{OH} + 2 \text{H} \xrightarrow{\text{AlCl}_3} \text{RC}_6\text{H}_4\text{OH} + \text{I}$, (cf. Smith, <i>loc. cit.</i>). Condensation of PhOH with MePhCHOH resulted chiefly in a mixt. of <i>o</i>- and <i>p</i>-MePhCHOH and a little of $\text{Et}_2\text{C}_6\text{H}_3\text{OH}$ as a result of the isomerization of the alk. radical, which could be effected either by the cleavage of H_2O from the alk. by the action of AlCl_3 or by the isomerization of alkylated phenol ether. The condensation of PhOH with primary alcs. was effected by the same method with 2 mols. of AlCl_3 to 1 mol. of PhOH and 1 mol. of alc. Condensation of MePhCHOH with PhOH in petr. ether at $110-20^\circ$ for 6 hrs. gave 52% <i>p</i>-$\text{HOC}_6\text{H}_4\text{CHMe}_2$ (III), m. 61°, (lit. ester m. $71-2^\circ$), and about 23% of nonphenolic fraction, b. $225-30^\circ$, contg. chiefly <i>p</i>-$\text{Me}_2\text{C}_6\text{H}_3\text{CHOC}_6\text{H}_4\text{CHMe}_2$ (IV). Anisole and MePhCHOH in petr. ether at 120° for 4 hrs. gave 38% of a mixt. of III and <i>o</i>-isomer, b. $215-17^\circ$, and 50% of <i>o</i>- and <i>p</i>-$\text{MeOC}_6\text{H}_4\text{CHMe}_2$ contaminated by a little IV. The condensation at $130-40^\circ$ for 5-6 hrs. resulted in 64% of <i>o</i>- and <i>p</i>-$\text{HOC}_6\text{H}_4\text{CHMe}_2$ and 30% $\text{MeOC}_6\text{H}_4\text{CHMe}_2$. <i>m</i>-Cresol and MePhCHOH heated at 125° for 6 hrs. gave 68% 3,4-$\text{HOC}_6\text{H}_3(\text{Me})_2$ (m. 112°), and 20% 3,4-$\text{Me}_2\text{C}_6\text{H}_3\text{CHOC}_6\text{H}_3(\text{Me})_2$ (b. $233-5^\circ$). PhOH and EtMeCHOH in ligroin</p> <p>see other side </p>	
<p>ASB-55A METALLURGICAL LITERATURE CLASSIFICATION</p> <p>19000 191000 192000 193000 194000 195000 196000 197000 198000 199000</p> <p>19000 191000 192000 193000 194000 195000 196000 197000 198000 199000</p>			

heated at 120-40° for 6 hrs. gave 52% of $\text{EtMeCHC}_6\text{H}_4\text{OH}$ (o-isomer, b. 229-33°, d₄ 0.9007, n_D 1.5182, M. R. 45.80; p-isomer, b. 235-7°, d₄ 0.9883, n_D 1.518), and 13% $\text{EtMeCHOC}_6\text{H}_4\text{CHMeEt}$. In the process of distn. the ether is given into the *tert*-butylphenol. Anisole and EtMeCHOH digested on a water bath for 2 hrs. gave 63%

$\text{MeOC}_6\text{H}_4\text{CHMeEt}$, b. 109-70°, d₄ 0.9413, n_D 1.503, M. R. 61.5, 16% $\text{MeOC}_6\text{H}_4(\text{CHMeEt})_2$, b. 140-2°, d₄ 0.9240, n_D 1.498, M. R. 90.72, and 13% $\text{HOC}_6\text{H}_4\text{CHMeEt}$. PhOH and MePrCHOH allowed to stand at room temp. for 24 hrs. and then heated at 40-50° for 4 hrs. gave 57.5% of phenolic and 11% of nonphenolic fraction. From the phenolic fraction crystd. out $\text{EtCHC}_6\text{H}_4\text{OH}$, m. 80° (Bz deriv., m. 40-1°); the residue on fractionation gave p- $\text{MePrCHC}_6\text{H}_4\text{OH}$, b. 150-6° (Bz deriv., b. 255°), and o-isomer, b. 140-50°, d₄ 0.9881, n_D 1.519 (Bz deriv., b. 340-4°). The nonphenolic fraction gave $\text{MePrCHC}_6\text{H}_4\text{OCHPrMe}$, b. 230-40°, and $(\text{MePrCH})_2\text{C}_6\text{H}_4\text{OCHPrMe}$, b. 245-60°, d₄ 0.900, n_D 1.4919. Anisole and MePrCHOH treated as above afforded $\text{MeOC}_6\text{H}_4\text{CHPrMe}$, b. 221-30°, d₄ 0.9284, n_D 1.490, and $\text{MeOC}_6\text{H}_4(\text{CHPrMe})_2$, b. 245-60°. The condensation of PhOH with MeOH could not be effected. PhOH and also ROH heated at

120-40° for 6 hrs. gave 62% phenolic and 11% nonphenolic products. The phenolic fraction consisted of 60% $\text{EtC}_6\text{H}_4\text{OH}$, b. 237°, m. 78-8.5°, and 40% o- $\text{EtC}_6\text{H}_4\text{OH}$, m. 137°, and the p-isomer, b. 216-20°. The nonphenolic fraction on redistn. was completely converted into $\text{EtC}_6\text{H}_4\text{OH}$. PhOH and PrOH gave 73.6% o- $\text{PrC}_6\text{H}_4\text{OH}$, b. 223°, d₄ 1.0008, and the p-isomer, b. 225-8°, and 17% $\text{PrC}_6\text{H}_4\text{OH}$, b. 240-5°; this on standing was converted into $\text{PrC}_6\text{H}_4\text{OH}$. PhOH and BuOH gave 72% phenolic and 14% nonphenolic products. o- $\text{BuC}_6\text{H}_4\text{OH}$, b. 233-5°, d₄ 0.9713; p-isomer, b. 240-60°, d₄ 0.9704 (Bz deriv., m. 25-6°). PhOH and *tert*- BuOH gave C_6H_5 and 82% $\text{MeC}_6\text{H}_4\text{OH}$, m. 98°. PhOH and *iso*- AmOH gave 3 amylphenols, of which only $\text{MeEtC}_6\text{H}_4\text{OH}$, m. 98-90°, was identified. About 25 references. Chas. Blanc

1ST AND 2ND SECTIONS

PROCESSING AND PROPERTIES INDEX

CONDENSATION OF ALCOHOLS WITH AROMATIC HYDROCARBONS IN THE PRESENCE OF ANHYDROUS ALUMINUM CHLORIDE. III. CONDENSATION OF PRIMARY ALCOHOLS WITH BENZENE AND TOLUENE. I. P. Luketevanik and G. Vukhrova. *J. Gen. Chem. (U.S.S.R.)* 7:632 6(1937); cf. C. A. 30, 412^a and preceding abstr.—The condensation of C_6H_5 and $PhMe$ with primary alcoh. proceeds more difficultly than with secondary alcoh., requiring a large excess of $AlCl_3$ (1.3-2 mols. for 1 mol. of alc.) and longer heating (4-10 hrs.) at higher temps. (110-40°). At optimum conditions 80-94% of mixed alkylated products and no unsubst. hydrocarbons are formed. Greater $AlCl_3$ concn. and higher temps. increase the formation of polyalkyl derivs. The alkylhydrocarbons are mixts. of *p*- and *m*-isomers with the latter as a chief reaction product (cf. Shoemith and McGeehan, C. A. 25, 281). $PhMe$ is generally more reactive. With $MeOH$ it gives products of various degrees of alkylation, while C_6H_5 fails to react. The 2-stage condensation (prepn. of Al alcoholate and heating it with C_6H_5 or $PhMe$) resulted in smaller yields of alkylated hydrocarbons. The nature of the diethylphenyl, $Rt(C_6H_5)_2Rt$ (I), $(C_6H_5)_3Rt$ (II) and diethyltriphenyl, $Rt(C_6H_5)_3Rt$ (III), formed from C_6H_5 and $EtOH$, is being investigated. Unless otherwise stated, the values given below after the b. p. are for d₄²⁰, n_D²⁰ and M. R., resp. Heating 16 g. (1/2 mol.) $EtOH$, 87 g. (9/2 mol.) $AlCl_3$ and 120 g. C_6H_5 at 120-30° for 10 hrs. resulted in 48.5% $PhRt$, b.p. 134-6°, 0.8722, 1.4957, 33.48 and 35% of polyalkylbenzenes. This redistd. gave *m*- C_6H_4Rt , b.p. 179-81°, 0.8636, 1.4978, 44.71; C_6H_5Rt , b.p. 215-17°, 0.8922, 1.5006, 54.03, mol. wt. 170 (calcd. 160); C_6H_5Rt , b. 240-50°, 0.8999, 1.5032, 63.28, mol. wt. 180 (calcd. 180); I, b. 290-311°, 0.938, 1.5425, 64.28, mol. wt. 334 (calcd. 210), and II m. 281°. From 50 g. (0.5 mol.) $PrOH$, 87 g. (9/2 mol.) $AlCl_3$ and 120 g. C_6H_5 at 110° for 10 hrs. resulted 51.5% $PhPr$, b. 160-8°, 0.866, 1.4911, 40.34 and 30.5% *m*- C_6H_4Pr , b. 215-18°, 0.9137, 1.5155, 53.89. Condensation of 23 g. (0.5 mol.) $EtOH$, 90 g. (0.75 mol.) $AlCl_3$ and 80 g. $PhMe$ at 140° for 8 hrs. gave 74% *m*- C_6H_4Et , b. 167-9°, 0.8992, 1.4937, 40.28 and the *p*-isomer, b. 160-3°, 0.8907, 1.4902, 40.23 and 20% *m*- C_6H_4Et , b. 160-3°, 0.8907, 1.4902, 40.23 and 20% *m*- C_6H_4Et . A mist. of (the position of the Rt groups was not detd.). A mist. of 30 g. (0.5 mol.) $PrOH$, 90 g. (0.7 mol.) $AlCl_3$ and 80 g. $PhMe$ heated at 125° for 4 hrs. afforded 6.5% *m*- C_6H_4Pr , b. 243-7° and 83% *m*- C_6H_4Pr , b. 182-5°, 0.8982, 1.4945, 44.84, and the *p*-isomer, b. 182-5°, 0.8982, 1.4945, 44.84. IV. Condensation of diphenyl alcohol with naphthalene. I. P. Luketevanik and G. Vukhrova. *J. Gen. Chem. (U.S.S.R.)* 7:632 6(1937); cf. C. A. 30, 412^a and preceding abstr.— $C_{10}H_8$ is easily alkylated by primary, secondary and tertiary alcoh., forming a mixture of mono-, di- and tri-alkylated products. The nature of the alkyl groups in the products is being investigated. The condensation of $C_{10}H_8$ with primary alcoh. is used at a ratio of 0.5 mol. to 1 mol. of tertiary alc. (excess $AlCl_3$ and prolonged heating result in poly-alkylphenylbenzenes), 1 mol. to 1 of secondary alc. and 1.5 : 1 alkylphenylbenzenes). The distribution of the position of alkyl groups in the prod. is based on the oxidation of alc.

see other side →

1930-1940 METALLURGICAL LITERATURE CLASSIFICATION

1930-1940 METALLURGICAL LITERATURE CLASSIFICATION

1930-1940 METALLURGICAL LITERATURE CLASSIFICATION

(CONTINUED)

CA Condensation of alcohols with aromatic hydrocarbons in the presence of anhydrous aluminum chloride. III. Condensation of primary alcohols with benzene and toluene. I.P.Tsukervanik and G. Vikhrova. J. Gen. Chem. (U.S.S.R.)

10

In the condensation of primary alcohols with aromatic hydrocarbons in the presence of anhydrous aluminum chloride, the reaction proceeds by intermediate cleavage of H_2O with the formation of cycloalkenes and cycloalkyl chlorides: $C_6H_5OH + AlCl_3 \rightarrow C_6H_5 + AlCl_2OH + HCl$; $C_6H_5 + HCl \xrightarrow{AlCl_3} C_6H_5Cl$; $C_6H_5 + C_6H_5 \xrightarrow{AlCl_3} Ph-C_6H_5$; $C_6H_5 + C_6H_5Cl \xrightarrow{AlCl_3} Ph-C_6H_4Cl + HCl$. C. B.

the condensation of secondary cyclic alcs. with aromatic hydrocarbons proceeds by intermediate cleavage of H_2O with the formation of cycloalkenes and cycloalkyl chlorides: $C_6H_5OH + AlCl_3 \rightarrow C_6H_5 + AlCl_2OH + HCl$; $C_6H_5 + HCl \xrightarrow{AlCl_3} C_6H_5Cl$; $C_6H_5 + C_6H_5 \xrightarrow{AlCl_3} Ph-C_6H_5$; $C_6H_5 + C_6H_5Cl \xrightarrow{AlCl_3} Ph-C_6H_4Cl + HCl$. C. B.

isomer (identified as $m-C_6H_4(CO_2Me)_2$, m. 63°) and $sym-C_6H_4(C_6H_5)_2$, m. 68° , identified as tri-Me trimesate, m. 144° . PhMe is less reactive, requiring more $AlCl_3$ and a large excess of PhMe. C_6H_5OH (70 g.), 80 g. $AlCl_3$ and 200 cc. PhMe gave 72% of a fraction, b.p. $254-60^\circ$, $d_4^{20} 0.9343$, $n_D^{20} 1.5342$, M. R. 57.05, consisting of p - and m - $MeC_6H_4C_6H_5$, identified as p - and m - $C_6H_4(CO_2Me)_2$, resp., and 18% $3,5$ - $Me_2C_6H_3(C_6H_5)_2$, m. 93.5° , identified as tri-Me trimesate. From the decomposition of C_6H_5OH with $AlCl_3$ with the formation of 14-20% C_6H_5 and 20-5% C_6H_5Cl it is assumed that by analogy with tertiary alcs.,

ASB-31A METALLURGICAL LITERATURE CLASSIFICATION

100 AND 200 GROUPS

PROCESSES AND PROPERTIES INDEX

10

Condensation of alcohols with aromatic hydrocarbons in the presence of anhydrous aluminum chloride. VI. Alkylation of halogenated aromatic hydrocarbons by alcohols. I. P. Tsukervanik. *J. Gen. Chem.* (U. S. S. R.) 8, 1512-15 (1938); *cf. C. A.* 31, 5789. The preliminary exptl. data on the condensation of excess of PhCl , $o\text{-ClC}_6\text{H}_4\text{Me}$ (I) and $1\text{-Cl}_2\text{C}_6\text{H}_3\text{Cl}$ (II) with alcs. in the presence of AlCl_3 at elevated temps. show that the alkylation of aryl halides proceeds analogously to that of unsubstituted aromatic hydrocarbons with the formation of corresponding monoalkyl- (40-65% yield) and polyalkylaryl chlorides. The alkyl derivs. represent mixts. of *p*- and some *m*-isomers, the constitution of which was detd. by oxidation to the corresponding chlorobenzoic acids. The condensation with primary alcs. (EtOH) proceeds most difficultly (requiring a large excess of AlCl_3 (1.5 g./mol.)); with secondary alcs. less difficultly (1 g./mol. AlCl_3); and with tertiary alcs. very easily (0.5 g./mol. AlCl_3). Greater AlCl_3 concn. and higher temps. increase the formation of polyalkyl derivs. The reactions are effected by gradual addn. of AlCl_3 to a mixt. of an aryl chloride and an alc. and refluxing on a boiling water bath for 2-3 hrs. Unless otherwise stated, the values given below are for b_p and d_4^{25} , resp. The reaction of 37 g. *tert*-BuOH and 100 g. PhCl with 30 g. AlCl_3 gave 11 g. polyalkyl chlorides (III) and 65% of isomeric *tert*-BuC₆H₄Cl; *p*-isomer, 210-12°, 1.0239, n_D^{25} 1.5125. PhCl (100 g.) and 44 g. *tert*-AmOH with 25 g. AlCl_3 gave 8 g. III and 50% *p*- and *m*-*tert*-AmC₆H₄Cl; *p*-isomer, 224-9°, 1.0008, n_D^{25} 1.5139. PhCl (100 g.) and 60 g. *iso*-PrOH with 120 g. AlCl_3 gave 62% chlorocumenes; *p*-isomer, 195-7°, 1.0022, n_D^{25} 1.514. PhCl (100 g.) and 44 g. *sec*-BuOH with 50 g. AlCl_3 gave 8 g. III and 50% isomeric *sec*-BuC₆H₄Cl; *p*-isomer, 212-14°, 1.001, n_D^{25} 1.4996. PhCl (100 g.) and 30 ml. EtOH with 120 g. AlCl_3 gave 10 g. III and 40% *p*-EtC₆H₄Cl, 179-80°, 1.0463, n_D^{25} 1.5235. PhCl (150 g.) and 74 g. *iso*-BuOH with 200 g. AlCl_3 gave 25 g. III and 30% of monoalkyl derivs., from which was isolated *p*-*tert*-BuC₆H₄Cl. PhCl (100 g.) and 44 g. *iso*-AmOH with 80 g. AlCl_3 gave 15 g. III and 35% monoalkyl derivs.; *p*-*tert*-AmC₆H₄Cl, 224-30°, 1.007, n_D^{25} 1.5139. I (25 g.) and 19 g. *tert*-BuOH with 15 g. AlCl_3 gave 45% monoalkyl derivs.; *tert*-BuMeC₆H₄Cl, 225-30°, 1.0070, n_D^{25} 1.5145. II (120 g.) and 44 g. *tert*-AmOH with 30 g. AlCl_3 gave 15 g. III and 60% monoalkyl derivs.; *tert*-amyl-1-chloronaphthalene, 310-12°, 1.1235, n_D^{25} 1.6009. II (75 g.) and 30 g. *iso*-PrOH with 60 g. AlCl_3 gave 12 g. III and 45% *iso*-propyl-1-chloronaphthalene, b. 305-10°, 1.1421, n_D^{25} 1.6043. Chas. Blanc

ASH-51A METALLURGICAL LITERATURE CLASSIFICATION

GROUP	SYMBOL	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100		


TSUKERVANIK, I. P.

Benzylstibine and its derivatives. I. P. Tsukervanik and D. Shadrinov. *J. Gen. Chem. (U. S. S. R.)* 7, 1527-31 (1937). — PhCH_2MgCl and SnCl_4 in a CO_2 atm. give *tri-benzylstibine* (I), m. 90° , which is easily oxidized by air, especially in soln., to *tribenzylstibine oxide* (II). I reacts with Cl_2 and Br_2 , or II with HCl and HBr , to give the *di-chloride* m. $100-1^\circ$, and the *di-bromide* m. $102-4^\circ$, resp. When I is heated with SnCl_4 , it forms $(\text{PhCH}_2)_2\text{SnCl}_2$, m. $187-8^\circ$, which gives with dil. NaOH soln. $[(\text{PhCH}_2)_2\text{SnO}]_2$. With HBr this forms $(\text{PhCH}_2)_2\text{SnBr}_2$, m. $150-1^\circ$ (decomps.). H. M. Leicester

Condensation of alcohols with aromatic compounds in the presence of aluminum chloride. IX. Condensation of 1,2-methylcyclohexanol, menthol and borneol with benzene and toluene. I. P. Tsukeryanik and N. G. Shkurva, *Gen. Chem.* (U.S.S.R.) 8, 1800 (1962) (in English, 1962) (1962); cf. C. A. 33, 4587. The following compounds were prepd.: *methylethylbenzene*, $C_{11}H_{14}$ (42.5% yield), b_p 142.3°, d_4^{20} 0.8308, n_D^{20} 1.5290; *methylethyltoluene* isomers (50% yield), b_p 152.4°, d_4^{20} 0.8291, n_D^{20} 1.5205; *menthylbenzene* $C_{11}H_{14}$ (51.5% yield), b_p 151.4°, d_4^{20} 0.8342, n_D^{20} 1.5207, *p*- and *m*-*menthyltoluene* (61% yield), b_p 181.8°, d_4^{20} 0.8244, n_D^{20} 1.5185; *bornylbenzene*, $C_{11}H_{14}$ (30% yield), b_p 165.8°, d_4^{20} 0.8535, n_D^{20} 1.5250; *p*- and *m*-*bornyltoluene* (64% yield), b_p 131.3°, d_4^{20} 0.8500, n_D^{20} 1.5275. In the prepn. of 1,3-menthene, b_p 165.0°, d_4^{20} 0.8103, n_D^{20} 1.1170, and *tert*-menthylchloride, $C_{11}H_{17}Cl$, b_p 117.8°, d_4^{20} 0.9105, n_D^{20} 1.4070, were the chief products when insufficient $AlCl_3$ was used. Small amts. of polyalkylated hydrocarbons were always obtained but not identified. John Livak

ASB.SLA METALLURGICAL LITERATURE CLASSIFICATION

1ST AND 2ND ORDERS										3RD AND 4TH ORDERS									
PROCESSING AND PROPERTIES INDEX																			
CA																			
<p>Reciprocal action of acetylketene and alcohols. 1. I. P. Tsukervanik and I. A. Ermolenko. <i>Bull. univ. Aie' centrale</i> 22: 215-20 (in German, 220) (1938).—The reaction between <i>acetylketene</i> and various alcs. and some other compds. has been investigated. The acetylation proceeds even in the cold in the absence of catalysts. The catalysts tested were quinoline, H_2SO_4, and $ZnCl_2$. Only in acetylating <i>glycerol</i> the presence of H_2SO_4 is of advantage. The reaction proceeds smoothly and the yield is almost theoretical in the case of primary and secondary alcs., whereas the acetylation of tertiary alcs. proceeds much more slowly. Glycerol and <i>glycol</i> give a mixt. of mono- and diacetates. Unsatd. alcs. react only with the HO group with the formation of normal acetates. The acetylation of <i>p-anisidine</i> gives 98% of pure acetate. <i>Aspirin</i> is obtained in 96% of the theoretical yield from <i>salicylic acid</i>. Gertrude Berend</p>																			
<p>ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION</p> <p>FROM: STIMULUS TO: BOMBAV TO: BOMBAV</p> <p>STIMULUS BOMBAV BOMBAV</p>																			



Condensation of aldehydes and ketones with aromatic hydrocarbons in the presence of anhydrous aluminum chloride. I. Condensation of aliphatic ketones with phenols. I. P. Tunkervanik and Z. N. Nazarova. *J. Gen. Chem.* (U. S. S. R.) 9, 33-5 (1939); cf. *C. A.* 31, 5778. The interaction of aliphatic ketones with PhOH in the presence of excess $AlCl_3$ (1.5 mols.) in the cold for 1-3 days yields dialkyl diphenylmethanes, which on heating decompose to give *p*-alkyl phenols. The latter are formed directly in 40% yield by condensation of the reaction mixt. on the water bath for 3-4 hrs. The mechanism of the condensation reaction is analogous to the similar procedure in the presence of fuming HCl studied by Diamm (*J. Russ. Phys.-Chem. Soc.* 23, 488, 523, 601 (1891); cf. *loc. cit.*). In this way were prepd. *isopropylphenol* (from Me_2CO and PhOH), b. 200-30° (and considerable resinification products) and *sec-amylphenol* (I) (from Et_2CO), m. 79°, b. 245-60° (*lit. deriv.*, m. 48-50°). Me_2CO also formed I, b. 245-50°, d₄²⁰ 0.97, n_D²⁰ 1.5215. It gives a *lit. deriv.*, b. 340-50°, *lit. deriv.*, b. 254-5°, and *Me ether*, b. 232-4°, which with Cl_2 is oxidized to anisic acid, m. 179°. Condensation of *m*-cresol with Me_2CO produced an anhydride, m. 131°, described by Zincke and Gaebel (*C. A.* 6, 1753). Chas. Blanc

ASD S.A. METALLURGICAL LITERATURE CLASSIFICATION

10

69

Alkylation of aromatic hydrocarbons by dihalogen derivatives. 1. Condensation of 1,3-chlorobromopropane with benzene. I. Tsukervanik and K. Yatsimirskii. *J. Gen. Chem.* (U.S.S.R.) 10, 1075 (1940). When $\text{Cl}(\text{CH}_2)_3\text{Br}$ and C_6H_6 react in the presence of AlCl_3 at $6-12^\circ$, 40% $\text{Ph}(\text{CH}_2)_3\text{Br}$ (I) is formed. At $80-5^\circ$, 60% $\text{Ph}(\text{CH}_2)_3\text{Ph}$ is formed, with some PhPr as a by-product, but no I is obtained. H. M. Leicester.

Lab. of Organic Chem., Am. Resin State Univ.

ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION

1ST AND 2ND CROSS										3RD AND 4TH CROSS									
PROCESS AND PROPERTIES INDEX																			
ca										10									
<p>The alkylation of aromatic compounds by alcohols in the presence of anhydrous ferric chloride. Z. N. Nazarov and I. P. Tsykervanik, <i>J. Gen. Chem.</i> (U. S. S. R.) 10, 1181-5(1946).—Tertiary alcs. react easily at room temp. with C_6H_5, $MePh$, C_6H_5, $PhOH$, <i>m</i>- and <i>p</i>-cresols and $MeOPh$ to give 70-85% yields of monoalkyl deriva., in the presence of 0.5 equiv. $FeCl_3$ per equiv. of alc. If the mixt. is heated, some <i>p</i>-dialkyl compds. are also formed. With secondary alcs., a 1:1 mixt. of $FeCl_3$ and alc. is required and the mixt. must be heated to 50-60° to get 38-78% yields of <i>o</i>- and <i>p</i>-derivs. With primary alcs., rate of reaction depends on structure, branched-chain alcs. reacting faster than normal alcs. H. M. Leicester</p>																			
<p>ASD-51A DETAILING LITERATURE CLASSIFICATION</p> <p>GROUP 1: 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20</p> <p>GROUP 2: 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20</p>																			

COMMON ELEMENTS		PROCESSES AND PROPERTIES INDEX		COMMON ELEMENTS	
C B		10			
<p>The condensation of aromatic compounds with acids.</p> <p>1. Condensation with hydrocarbons, phenol and phenols. I. Tsukervanik and I. Terent'eva. <i>J. Gen. Chem.</i> (U. S. S. R.) 10, 1103-7 (1940).—<i>iso</i>-BuCO₂H (30 g.) in 100 cc. PhMe was treated with small portions of AlCl₃ (total 75 g.). After adding all the AlCl₃, the reaction mixt. was allowed to stand for 12 hrs. and then was heated in a water bath for 3-6 hrs. <i>p</i>-Methylisovalerophenone, obtained in 72% yield, b_m 224-5°, b₁₀ 137°, d₄ 0.9574, n_D 1.5085, MR_D 54.39; semicarbazone, m. 215-16°; oxime, 65°. Oxidation of the product with 10 parts of 5% HNO₃ in a sealed tube for 10 hrs. yielded <i>p</i>-McC₂H₄CO₂H. Condensation of PrCO₂H in PhMe in the presence of AlCl₃ under the same conditions yielded <i>p</i>-methylbutyrophenone (72.5%), b. 248-50°, d₄ 0.9000, n_D 1.5232, MR_D 51.20; semicarbazone, m. 200-10°. On the oxidation of the product <i>p</i>-McC₂H₄CO₂H was reph. Under the same conditions EtOPh condensed with <i>iso</i>-BuCO₂H in the presence of AlCl₃, yielding <i>p</i>-ethoxyisovalerophenone (82%), b_m 153-7°, d₄ 1.0406, n_D 1.5533, MR_D 61.01 (semicarbazone, m. 191-2°; oxime, m. 118-19°); and <i>p</i>-hydroxyisovalerophenone (9%) m. 95-6°. Phenole with PrCO₂H yielded <i>p</i>-ethoxybutyrophenone (76%), b. 129°, n_D 1.5390, MR_D 57.03 (semicarbazone, m. 181°; oxime, m. 103-4°), and <i>p</i>-hydroxybutyrophenone, m. 91-2° (semicarbazone, m. 167-9°; oxime, m. 83-4°). The condensation of PhOH with <i>iso</i>-BuCO₂H in C₆H₆ under the same conditions yielded <i>p</i>- and <i>o</i>-hydroxyisovalerophenones (total yield 63%); the <i>o</i>-deriv. b. 248-50°, d₄ 1.0197, n_D 1.5126, MR_D 52.99. A. A. Podgorny</p>					
<p>ABB-SLA METALLURGICAL LITERATURE CLASSIFICATION</p> <p>EXHIBIT 10000</p> <p>EXHIBIT 10000</p>					

Condensation of alcohols with aromatic hydrocarbons in the presence of aluminum chloride. Condensation of cycloheptanol with benzene and toluene. N. G. Sidorenko and I. E. Tsukerman. *J. Gen. Chem.* (U.S.S.R.) 10, 2073-6 (1940). - A yield of 46-50% of cycloheptylbenzene (I) and cycloheptyltoluene (II) was obtained on condensation of cycloheptanol (III) with benzene (IV) and toluene (V). III was prepd. by the reduction of cycloheptanone by the method of Mosettig and Burger (C. I. 44, 6767); 10 g. of III, 12 g. $AlCl_3$ and 100 ml. IV yielded 7 g. I, b.p. 132-3°, d_4^{20} 0.9410, n_D^{20} 1.5217, M_R 56.42, calcd. 56.43. Oxidizing I with 15% HNO_3 in sealed tubes at 140-160° yielded IIaOH, m. 119°. *p*-Cycloheptylnitrobenzene (VI) was obtained on nitrating I with a nitration mixt. VI b.p. 283-10°. Oxidizing VI with 15% HNO_3 in sealed tubes gave *p*- $O_2NC_7H_{13}CO_2H$, m. 280°. The *p*-amino deriv. (VII) of I was obtained on reduction of VI with Sn and HCl, and was purified by steam distn. The benzoate of VII (from alc.) m. 173°; acetate (from deriv. of phenylcyclohexylmethane (VIII) and 1-methyl-2-phenylcyclohexane (IX), which were specially prepd., differed from the derivs. of I. **Synthesis of VIII.** Phenylcyclohexylcarbinol (X) was obtained by Grignard's reaction from 10 g. Mg, 68 g. bromocyclohexane, 2.4 g. NaH in 100 ml. of abs. ether. The yield was 34.5 g. (45.5%), b.p. 130-42°. Ph cyclohexyl ketone (XI) was formed by the oxidation of X. To 31.5 g. of the heated X was added dropwise 18.5 g. $Na_2Cr_2O_7$ in 45 ml. of H_2O and 13

ml. H_2SO_4 . The reaction product was acid with ether. After the ether was evapd. thin needles of XI, m. 54°, were obtained. The yield was 31 g. (91%). Phenylcyclohexylmethane (XII) was obtained by the reduction of XI by the Clemmensen method. The yield was 32%, b.p. 137°, d_4^{20} 0.9436, n_D^{20} 1.5255, M_R 56.58, calcd. 56.49. The (nitrophenyl)cyclohexyl ethane, b.p. 18°, with Sn and HCl. The *N*-H₂ deriv. (from dil. alc.) m. 161-2°. The *N*-Ac deriv. (from dil. alc.) m. 170-1°. **Synthesis of IX.** 3-Methyl-2-phenylcyclohexene (XIII) was obtained by Grignard's reaction from 1 g. Mg, 20 g. bromobenzene and 18 g. of 2-methylcyclohexanone in 50 ml. of abs. ether. The yield was 12 g. (42%), b.p. 112-15°. 1-Methyl-2-phenylcyclohexane (XIV) was obtained on heating 12 g. of XIII with 12.5 ml. H_2 (of 1.7) and 0.75 g. of red P in a sealed tube, for 6 hrs., at 160-200°. XIV b.p. 91-2°, d_4^{20} 0.9599, n_D^{20} 1.5300, M_R 56.75, calcd. 56.63. The nitro and the corresponding amino compds. were obtained from XIV by the usual method. The *N*-H₂ deriv., m. 169-74°. The *N*-Ac deriv. was a liquid, b.p. 195-200°. It was obtained on treating 5 g. of III with 50 ml. of V and 6 g. of $AlCl_3$. The yield was 4 g. of II (a mixt. of *m*- and *p*-isomers), b.p. 110-12°, d_4^{20} 0.9350, n_D^{20} 1.5255, M_R 60.81, calcd. 61.05. It oxidized with dil. HNO_3 yielded a mixt. of *tert*- and isophthalic acids, which were obtained by their *di*-Me esters. The conclusion is drawn that the condensation of III with IV and V proceeds only to a small extent.

James J. Lachin

ca

10

Alkylation of aromatic compounds with alcohols in the presence of anhydrous FeCl_3 . Z. N. Nazarova and I. P. Tsubernanik, *J. Gen. Chem.* (U. S. S. R.) 14, 77 (1944) (Russian summary).--Attempts to effect a condensation between benzene and PhOH on one hand and primary alcs. (EtOH , PrOH and BuOH) on the other hand failed to give more than traces of alkaryl derivs. PhCH_2OH (20 g.), benzene (50 g.) and 32 g. FeCl_3 heated to 70° for 4 hrs. gave 58% Ph_2CH_2 , b. $200-5^\circ$, m. 27° . PhCH_2OH (20 g.), PhOH (20 g.) and 15 g. FeCl_3 mixed, let stand for 24 hrs. and heated on a steam bath for 2 hrs. gave 58% *benzylphenol*, b. $300-30^\circ$, m. 83° . Allyl alc. (1A g.), 40 g. benzene and 25 g. FeCl_3 were mixed with cooling, let stand for 24 hrs. and finally heated for 2-3 hrs. on a steam bath to 80° to yield 8% crude *allylbenzene* and 32% (*2-chloropropyl*)benzene; the purified compds. had the following consts., resp.: b. $154-6^\circ$, n_D^{20} 1.5110, d_4^{20} 0.8660, and b.m. $200-5^\circ$, n_D^{20} 1.5147, d_4^{20} 1.0249. This reaction conducted at room temp. (3 days' standing) gave the 2 products in reverse ratio: 32% and 8%, resp. Allylbenzene yields a *tribromide*, m. 118° . Similarly, 1A. 3Ae gave *allyloluene*, b. $178-80^\circ$, n_D^{20} 1.6085, d_4^{20} 0.8648, and (*2-chloropropyl*)luene, b. $220-30^\circ$, n_D^{20} 1.5194, d_4^{20} 0.8634. G. M. Kosolapoff

ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION

Ca

10

The mechanism of alkylation reactions under the influence of anhydrous FeCl_3 . Z. N. Nazarova and I. P. Tsukervanik. *J. Gen. Chem. (U.S.S.R.)* 14, 236-44 (1944) (English summary); cf. *C.A.* 39, 916¹.— EtBr and FeCl_3 form EtBr and EtCl , PrBr gives iso-PrCl , and iso-AmBr gives C_6H_{10} and mixts. of AmCl and AmBr . When EtBr and C_6H_6 are heated with FeCl_3 , HCl , EtCl , and PhBr are formed. BuI gives an analogous reaction. Under similar conditions PrBr gives iso-PrCl , PhBr , $\text{Br-C}_6\text{H}_4\text{Pr}$, and a little PhPr ; iso-BuBr gives PhBr , tert-BuPh , iso-BuPh and a mixt. which probably contains $\text{Br-C}_6\text{H}_4\text{Bu}$ and $\text{Bu-C}_6\text{H}_4$. BuCl and AmCl give almost no reaction with C_6H_6 and FeCl_3 . EtOH and FeCl_3 give 87% EtCl . With FeCl_3 , PrOH gives iso-PrCl and a little PrCl . BuOH gives BuCl and some sec-BuCl . iso-BuOH gives tert-BuCl and diisobutylene. FeCl_3 dissolves in iso-AmOH , and on standing very hygroscopic crystals sep. which m. 48° , evolve HCl at 58° , and solidify at 105° to a green, nonhomogeneous mass. iso-PrOH gives iso-PrCl ; tert-BuOH gives C_6H_6 , sec-BuCl , and tert-BuCl ; tert-AmOH gives C_6H_{10} and tert-AmCl . iso-PrOPh is isomerized by FeCl_3 to $\text{PrC}_6\text{H}_4\text{OH}$ and much tar is formed. These reactions are interpreted in terms of formation of an ionized complex of the type $(\text{ROH} \cdot \text{FeCl}_3)^+\text{Cl}^-$ which decomps. in various ways to give the products observed. Alkylation with FeCl_3 is thus not a simple condensation, as with AlCl_3 , but involves rupture of C bonds and deep-seated changes in structure. H. M. Leicester

Lab. Organic Chem. University, State Univ., -1943-

454-55A METALLURGICAL LITERATURE CLASSIFICATION

FROM DIVISION

REPORT ONE ONLY

<p>ca</p> <p>ASH-31A METALLURGICAL LITE</p>	<p>Condensation of alcohols with aromatic compounds in the presence of aluminum chloride. XI. Mechanism of alkylation of aromatic hydrocarbons by alcohols. I. P. Fokhtyaynik (Middle Asiatic State Univ.). J. Gen. Chem. (U.S.S.R.) 15, 635-8(1945) (English summary).-- The cleavage of reaction products of alcs. with AlCl₃ was studied at temps. above 100°; contrary to Norris and Sturgis (C.A. 33, 6231), only low yields of alkyl chlorides were obtained. It was shown that addn. of 5-10% FeCl₃ to AlCl₃ markedly raises the yield of RCl on cleavage of the AlCl₃OR. PrOH (23 g.) was added over 5 hrs. to 67 g. AlCl₃; vigorous evolution of HCl was observed and the resulting solid was slowly heated to 135° with collection of the distillate in a well-cooled receiver; there were obtained 37% EtCl and 3 g. Et₂O. Similarly, iso-PrOH gave 25% iso-PrCl and PrOH gave 10% PrCl (at 115-20°). EtOH (23 g.) was similarly added to 64 g. AlCl₃ and 3.5 g. FeCl₃; the resulting mass m. 100°, while heating to 125° produced foaming and evolution of EtCl, with the reaction being complete at 150°; there was obtained 70% EtCl. Use of 60 g. AlCl₃ and 7 g. FeCl₃ as above gave 96% EtCl. Iso-PrOH (12 g.), 20 g. AlCl₃, and 3 g. FeCl₃ gave 73% iso-PrCl; 30 g. PrOH, 60 g. AlCl₃, and 7 g. FeCl₃ gave 54% PrCl, while 37 g. BuOH, 60 g. AlCl₃, and 7 g. FeCl₃ gave 10% BuCl. Iso-AmOH failed to yield iso-AmCl. AlCl₃ (30 g.) was slowly added to 20 cc. iso-BuOH in 50 cc. CS₂ with cooling, after which the mass was warmed on a water bath until the HCl evolution ceased and 2 layers formed; the top layer on cooling gave a crop of colorless crystals of iso-BuOAlCl₂ (22.5%); use of petr. ether instead of CS₂ gave an analogous result with a 32% yield of the product; treatment of this with water gave iso-BuOH, while addn. of 20 g. of the complex to 30 cc. benzene and 20 cc. AcCl gave 60% PhAc (the reaction was run with effective cooling); similar reaction with BzCl gave 50% PhCO₂Et. S. M. Kosolapoff</p>	<p>10</p> <p>MBR., LAB. ORGANIC CHEMISTRY, SVERDLOV STATE UNIV., -1940-</p>
---	---	---

1ST AND 2ND ORDERS										3RD AND 4TH ORDERS									
PROCESSING AND PROPERTY INDEX																			
10										10									
<p>Alkylation of aromatic compounds in the presence of phosphoric acid. I. Condensation of alcohols with aromatic hydrocarbons and halides. I. P. Trukervanik (Middle Asiatic State Univ.). <i>J. Gen. Chem.</i> (U.S.S.R.) 15, 599-703 (1945) (English summary).—It was shown that H_3PO_4 may be used for alkylation of aromatic compds. by alc., with yields up to 90-95% being obtained in the best cases. Considerable poly-substitution occurs, <i>p</i>-isomers predominating in disubstituted products, with occasional traces of the <i>o</i>-isomers. The iso alcs. usually isomerize into tertiary radicals. The mixt. of aromatic compd. and H_3PO_4 was treated, with stirring, with the alc. which was added slowly, and the mixt. was heated and stirred for 6-10 hrs.; on cooling, the org. layers were fractionated. Thus, 100 cc. benzene, 37 g. iso-BuOH, and 110 cc. H_3PO_4 (d. 1.85) gave, after 10 hrs. at 90°, 3 g. PhH, b. 165-70°, with almost all benzene being recoverable; the use of toluene and iso-AmOH as above gave 30% mixed amyltoluenes, principally <i>p</i>-AmC₆H₄Me, b. 204-6°. Toluene and iso-BuOH gave 50% mixed butyltoluenes, principally <i>p</i>-<i>tert</i>-BuC₆H₄Me, b. 193-6°. Toluene and iso-PrOH gave 80% <i>p</i>-cymene; toluene and EtOH gave a very low yield of ethyltoluene, b. 157-60°. Toluene and allyl alc. gave 10% allyltoluene, b. 183-6°, and 40% diallyltoluene, b. 302-7°. Toluene (75 cc.), 46 g. BuCl, and 110 cc. H_3PO_4, after 12 hrs. at 90°, gave 15% butyltoluene isomers, b. 185-95°, while <i>tert</i>-BuBr gave 37% <i>p</i>-Me₂CC₆H₄Me. PhBr (80 g.), 37 g. iso-BuOH, and 100 cc. H_3PO_4 gave, after 8 hrs. at 120-30°, 30% BuC₆H₄Br; a higher temp. gave a higher yield but also caused much tar formation; the use of PhCl gave 30% <i>p</i>-Me₂CC₆H₄Cl, b. 210-12°, while PhCl and <i>n</i>-PrOH gave 90% mixed propylchlorobenzenes, b. 185-95°, with a considerable amt. of <i>p</i>-PrC₆H₄Cl, b. 190-2°. Benzene and borneol at 70° (10 hrs.) gave 25% bornylbenzene, b. 145-8°.</p> <p>G. M. Kasolapoff</p> <p>mbr., LAB. ORGANIC CHEM., MIDDLE ASIATIC STATE UNIV.,</p>																			
<p>ASB-31A METALLURGICAL LITERATURE CLASSIFICATION</p> <p>FROM SYMBLIV</p> <p>FROM SYMBLIV</p> <p>FROM SYMBLIV</p>																			

10

Alkylation of aromatic compounds in the presence of phosphoric acid. II. Alkylation of phenol and anisole by phosphoric acid. V. Tambovtseva and I. Tankryan. *Gen. Chem. (U.S.S.R.)* 15, 820-1 (1940); *Ch. U.S.S.R.* 40, 3707. - PhOH and PhOMe were condensed in the presence of 1.5 mols. 95% H_3PO_4 with *tert*-BuOH at 70-80°, with *iso*-PrOH and *iso*-BuOH at 80-100°, and with PhOH and BuOH at 100-30° to form 80-95% of the corresponding *p*-alkyl derivs. and sometimes a small amt. of the *o*-derivs. Below these temps., reaction did not occur. During the condensation, primary groups isomerized to secondary, and secondary to tertiary. Isolation of $PhO(CH_2)_3OH$ from one reaction mixt. indicates that the reaction goes through the formation of acid esters of H_3PO_4 , which are in turn the alkylating agents. With PhOMe both *Pr* and *iso*-*Pr* alcs. yielded *p*-isopropylanisole, *b_m* 212-14°, *d₄* 0.9118, *n_D* 1.5070, and *o*-isopropylanisole, *b_m* 198-200°, *d₄* 0.9540, *n_D* 1.5030. Similarly, *Pr* and *iso*-*Pr* alcs. and phenol yielded the corresponding *p*-*iso*-*Pr* deriv., *b_m* 213-14°, *d₄* 1.0157, *n_D* 1.5092, and *o*-*iso*-*Pr* deriv., crystals, *m.* 61°. Other compds. prepd. were *p*-*tert*-butylanisole, *b_m* 211-15°, *d₄* 0.9135, *n_D* 1.5060, and *p*-*tert*-butylphenol, crystals, *m.* 99° (from *iso*-BuOH), and *p*-*sec*-butylanisole, *b_m* 198-10°, *d₄* 0.9110, *n_D* 1.5030 (from BuOH). H. M. Leicester

Mr., Lab. Organic Chemistry, Middle Aetiohic State Univ., - 1944 -

ASH-SLA METALLURGICAL LITERATURE CLASSIFICATION

10

Activity of halogen in organic compounds in Friedel-Crafts syntheses. I. Condensation of ethyl esters of aliphatic halogen-substituted acids with benzene. I. P. Tsukervanik and I. V. Terent'eva. *Doklady Akad. Nauk S.S.S.R.* 90, 257-60 (1945).—The position of Cl determines the nature of the products from the Friedel-Crafts reaction of C_6H_6 with esters of halo aliphatic acids; the activity of Cl rises with increased distance from the COO group. C_6H_6 (100 ml.), 40.5 g. EtO_2CCH_2Cl , and 68 g. $AlCl_3$, after standing 24 hrs., boiling 1.5 hrs., and heating 5 hrs. on a steam bath, gave 70% mixed hydrocarbons from which $EtPh$, a mixt. of isomeric $Et_2C_6H_5$, and 1,2,4- $Et_3C_6H_3$, were isolated: the latter gave 1,2,4- $C_6H_3(CO_2H)_3$, m. 215-18°, after oxidation by HNO_3 at 165-70°. C_6H_6 (100 ml.), 34 g. $EtO_2CCH_2CH_2Cl$, and 70 g. $AlCl_3$, after 24 hrs. standing and 6 hrs. at 80°, gave 68% $PhCH_2CH_2CO_2H$ (m. 40°, b. 280°), 7 g. $EtPh$, and 12 g. $PhCOCH_2CH_2Cl$ (b. 255 CA°, n_D^{20} 1.4832, d_4^{20} 1.0861); merely letting the mixt. stand 14 days gave only 74% $PhCH_2CH_2CO_2H$ and $EtPh$. C_6H_6 (50 ml.), 9 g. $EtO_2C(CH_2)_3Cl$, and 20 g. $AlCl_3$, let stand 36 hrs., kept 12 hrs. at 60-70°, and boiled 3 hrs. gave 2 g. $EtPh$, 4 g. $Ph(CH_2)_3CO_2H$, m. 51°, and 4.5 g. 3,4-dihydro-1(2H)-naphthalenone, b.p. 132-4°, n_D^{20} 1.5005, d_4^{20} 1.0905. G. M. Kosolapoff

ASM-51A METALLURGICAL LITERATURE CLASSIFICATION

197 AND 198 ORDERS

THE NEW YORK PUBLIC LIBRARY
ASTOR LENOX TILDEN FOUNDATION
JAN 10 1965

Pd

10

Thermal splitting of opianic acid. I. P. Tsukervanik and A. S. Sultanov (Middle-Asiatic State Univ.). J. Gen. Chem. (U.S.S.R.) 16, 1715-17(1946).—The reaction was studied as a source of veratrinaldehyde. Narcotine (tech., 300 g.) was added to 3 l. 5% H₂SO₄, heated to 60° and the mixt. heated to gentle boiling and treated slowly with 500 g. NaO over 1.5 hrs., with concurrent addn. of 1.5 l. 10% H₂SO₄. The mixt. was boiled 5 hrs., treated with a sat'd sol'n. of 25–30 g. KMnO₄, boiled 0.5 hr., and treated with 5 g. charcoal. After filtration and cooling 1.5 days, 63% opianic acid, m. 144–5°, was obtained. This was purified by soln. of 50 g. in 120 cc. 20% NH₄OH,

filtration, and treatment with 40 g. CaCl₂; after heating until soln., cooling, clarifying with charcoal, and filtration, pure opianic acid, m. 148°, was obtained on cooling and acidification with 10% HCl; 5 g. subjected to dry distn. gave 2 g. distillate which, after soln. in CCl₄, washing with 5% NaOH, and evapng., gave 7.5% veratrinaldehyde, isolated as the semicarbazone, m. 175°; repetition of the distn. at 170° and 10–12 mm. gave 4 g. distillate of the distn. contg. 0.6 g. veratrinaldehyde and 0.2 g. isovanillin. The results were improved by conducting the pyrolysis in the presence of Cu (Kahlbaum Cu bronze powder) in a small autoclave at 200–10° (20–5 atm. pressure) in benzene soln. The best result was obtained as follows: 50 g. opianic acid, 100 g. Cu, and 130 cc. CCl₄ were heated to 200–10° 5 hrs., after which the mass was continuously extd. with CCl₄ and veratrinaldehyde (bp. n 135–42°) was fractionally distd. from the ext. in 60% yield. G.M.K.

Mex., Lab. Organic Chemistry,
Middle Asiatic State Univ.
-1945-

COMMON ELEMENTS

METALLURGY

MINERALOGY

CHEMISTRY

PHYSICS

BIOLOGY

MEDICINE

LAW

HISTORY

SOCIAL SCIENCES

ARTS

GENERAL REFERENCE

SERIES

VOLUME

PAGE

DATE

TIME

BY

FOR

REMARKS

SIGNATURE

INITIALS

NUMBER

CLASSIFICATION

SUBJECT

AUTHOR

TITLE

PUBLICATION DATE

COUNTRY OF ORIGIN

LANGUAGE

FORMER LOCATION

ACQUISITION METHOD

PRICE PAID

DONOR NAME

DOLLAR AMOUNT \$

CENTS

TOTAL DOLLARS

FUND NUMBER

PROJECT NUMBER

GRANT NUMBER

AGENCY

PROGRAM

PILOT PROJECT

CONTRACT NO.

MODIFIED BY

DATE MODIFIED

APPROVED BY

DATE APPROVED

STATUS

COMPLETION DATE

PERCENTAGE COMPLETE

ACTUAL COST

ESTIMATED COST

VARIANCE

REASON FOR VARIANCE

ACTION REQUIRED

RESPONSIBLE PARTY

CONTACT INFORMATION

TELEPHONE

ADDRESS

ELECTRONIC MAIL ADDRESS

WEBSITE URL

OTHER CONTACT INFO

NOTES

ADDITIONAL COMMENTS

APPENDIX

REFERENCES

BIBLIOGRAPHY

PATENT LITERATURE

STANDARD SPECIFICATIONS

TEST METHODS

CALCULATIONS

RESULTS SUMMARY

CONCLUSIONS

RECOMMENDATIONS

FINAL REPORT

EXECUTIVE SUMMARY

INTRODUCTION

OBJECTIVES

SCOPE

DEFINITIONS

ABBREVIATIONS

SYMBOLS

UNITS

CONVERSION FACTORS

GLOSSARY

INDEX

TABLE OF CONTENTS

LIST OF FIGURES

FIGURE CAPTIONS

EXPLANATORY NOTES ON FIGURES

ACKNOWLEDGMENTS

DEDICATION

DISCLAIMER

COPYRIGHT NOTICE

TRADE MARK NOTICES

REGISTERED TRADE NAMES

PATENT REFERENCES

LEGAL OPINION

CERTIFICATE OF AUTHORITY

STATEMENT OF WORKING STATUS

DECLARATION OF INVENTION

ABSTRACT

KEYWORDS

THESIS STATEMENT

SUMMARY OF FINDINGS

CONTRIBUTION TO KNOWLEDGE

IMPlications FOR FUTURE RESEARCH

LIMITATIONS OF STUDY

STRENGTHS OF STUDY

WEAKNESSES OF STUDY

OPPORTUNITIES FOR IMPROVEMENT

CHALLENGES FACED

LESSONS LEARNED

BEST PRACTICES IDENTIFIED

RECOMMENDED ACTIONS

NEXT STEPS

LONG TERM GOALS

SHORT TERM GOALS

TIMELINE

MILESTONES

DELIVERABLES

RISKS ASSESSMENT

MITIGATION STRATEGIES

CONTINGENCY PLANS

MEL'KANOVITSKAYA, S.G.; TSUKERVANIK, I.P.

Allylation of phenol compounds in the presence of a copper catalyst. Dokl. AN Uz. SSR no. 11:40-44 '59. (MIRA 13:4)

1. Institut khimii AN UzSSR. 2. Chlen-korr. AN UzSSR (for Tsykervanik).
(Propene) (Eugenol)

SOV/7)-28-3-7/66

AUTHORS:

Mel'kanovitskaya, S. G., Tsukervanik, I. P.

TITLE:

Radical and Ionic Alkylation of the Aromatic Nucleus
(Radikal'noye i ionnoye alkilirovaniye aromaticeskogo
yadra) VII. Butylation of Benzene, Naphthalene, Phenol, and
Anisole (VII. Butilirovaniye benzola, naftalina, fenola i
anizola)

SYNOPSIS:

Zhurnal obshchey khimii, 1958, Vol. 28, Nr 8, pp. 2032-2038
(USSR)

ABSTRACT:

In connection with a previous paper (Ref 1) this paper at-
tempts to explain the peculiarity of the alkylation of dif-
ferent compounds with n-butyl chloride in the presence of
copper. As an example the free radical butylation of anisole
was investigated. n-Butylphenyltriazenes were used as the
source of free butyl radicals. In contrast to the benzylation
and the reaction with diphenylchloromethane the butylation
with copper proceeds much more difficultly (at 200-250°).
In reacting n-butylchloride with benzene, naphthalene, phenol,
and anisole the corresponding monobutyl and dibutyl aromatic
compounds resulted, and in the case of phenol a butyl ether

Card 1/3

Radical and Ionic Alkylation of the Aromatic Nucleus. SOV/79-28-8-7/66
VII. Butylation of Benzene, Naphthalene, Phenol and Anisole

of the butylphenol also resulted. Side products could not be determined. In every case it was the secondary butyl derivative which was produced. The proof of the structures was carried out by perbrominating and oxidizing. It was found that the alkylation product yield in the above reactions was 38-87 %. Thermal butylation also yielded alkyl derivatives. Under the experimental conditions used an isomerization of the primary butyl group to the secondary form always took place. The butylation of phenol follows the mechanism of indirect nuclear alkylation. Free butyl radicals produced by the thermal decomposition of n-butylphenyltriazene can be used to alkylate the anisole nucleus. There are 4 tables and 25 references, 11 of which are Soviet.

ASSOCIATION: Institut khimii Akademii nauk UzSSR (Institute of Chemistry AS UzSSR)

SUBMITTED: July 12, 1957

Card 2/3

Radical and Ionic Alkylation of the Aromatic Nucleus. RCV/79-20-8-7/66
VII. Bu~~l~~kylation of Benzene, Naphthalene, Phenol, and Anisole

Card 3/3

AUTHORS: Tsukervanik, I. P., Belinson, Z. Ya. SOV/79-28-8-8/66

TITLE: Condensations of Benzotrichloride With Benzene in the Presence of Aluminum Chloride (Kondensatsii benzotrikhlorda s benzolom v prisutstvii khloristogo alyuminiya)

PERIODICAL: Zhurnal obshchey khimii, 1958, Vol. 28, Nr 8, pp. 2038-2042 (USSR)

ABSTRACT: The authors investigated the condensations of benzotrichloride with benzene in order to produce the various substitutions of the chloride atom and to ascertain the conditions under which triphenylmethane and 9-phenylfluorene form. It was found that the amount of $AlCl_3$ present is a definite factor in this synthesis. Using 0,1 mole $AlCl_3$ (to 1 mole benzotrichloride), and independent of the temperature ($15-90^\circ$) and the reagent concentration, a yield of 70 % diphenyldichloromethane was obtained with a small side product of triphenylchloromethane. With 1 mole of $AlCl_3$, triphenylchloromethane was produced as the main product with a yield of 80-98 %. Further increases in the amount of $AlCl_3$ caused a hardening reaction and a decrease in the triphenylchloromethane yield. Other factors in this synthesis were studied, among them

Card 1/2